

On the Estimates of the Density of the Purely Discontinuous Girsanov Transforms of α -Stable-like Processes

Chunlin Wang
Department of Mathematics
University of Illinois
Urbana, IL 61801, USA
Email: cwang13@uiuc.edu

Abstract

In this paper, we study the purely discontinuous Girsanov transforms which were discussed in Chen and Song [3] and Song [9]. We show that the transition density of any purely discontinuous Girsanov transform of a α -stable-like process, which can be nonsymmetric, is comparable to the transition density of the α -stable-like process.

AMS 2000 Mathematics Subject Classification: Primary 60J45, 60J40; Secondary 35J10, 47J20.

Keywords and phrases: Transition density, Girsanov transform, martingale additive functional, α -stable-like processes, Brownian motion

Running Title: Density estimates of the purely discontinuous Girsanov transformed processes

1 Introduction

Suppose that $(X, \mathbb{P}_x)$ is a Brownian motion in $\mathbb{R}^d$ and $\mathcal{M}_t$ is the σ -field generated by $\{X_s, s \leq t\}$. Define

$$L_t = \exp \left(\int_0^t b(X_s) dX_s - \frac{1}{2} \int_0^t |b(X_s)|^2 ds \right),$$

where b is an $\mathbb{R}^d$ -valued function on $\mathbb{R}^d$. When b satisfies certain natural conditions, L_t is a nonnegative local martingale under each $\mathbb{P}_x$ and thus is a supermartingale multiplicative additive functional of X . L_t defines a family of probability measure $\{\tilde{\mathbb{P}}_x : x \in \mathbb{R}^d\}$ on $\mathcal{M}_\infty$ by $d\tilde{\mathbb{P}}_x = L_t d\mathbb{P}_x$ on $\mathcal{M}_t$. Under this family of measures $\{\tilde{\mathbb{P}}_x : x \in \mathbb{R}^d\}$, X_t is denoted by $\tilde{X}_t$. The process $\tilde{X}$ is called a Girsanov transform of X and the transition density of $\tilde{X}$ has Gaussian upper and lower estimates for certain natural function b .

The main purpose of this paper is to study upper and lower estimates on the transition densities of Girsanov transforms of α -stable-like processes (see definition 1.1).

Let's first recall the purely discontinuous Girsanov transforms studied in [3]. For convenience, we consider the state space $\mathbb{R}^d$ instead of a Lusin space E in [3]. Let $\mathcal{B}(\mathbb{R}^d)$ be the Borel σ -field and m a σ -finite measure on $\mathcal{B}(\mathbb{R}^d)$. Let $X = (X_t, \mathbb{P}_t)$ be a m -Hunt process on $\mathbb{R}^d$ and has a Lévy system (N, H) . That is for any nonnegative function f on $\mathbb{R}^d \times \mathbb{R}^d$ vanishing on the diagonal

$$\mathbb{E}_x \left(\sum_{s \leq t} f(X_{s-}, X_s) \right) = \mathbb{E}_x \left(\int_0^t \int_{\mathbb{R}^d} f(X_s, y) N(X_s, dy) dH_s \right),$$

for every $x \in \mathbb{R}^d$ and $t > 0$.

Let F be a bounded function on $\mathbb{R}^d \times \mathbb{R}^d$ that vanishes on the diagonal. We say $F \in \mathbf{J}(X)$ if

$$\lim_{t \downarrow 0} \sup_{x \in \mathbb{R}^d} \mathbb{E}_x \left(\int_0^t \int_{\mathbb{R}^d} |F|(X_s, y) N(X_s, dy) dH_s \right) = 0,$$

see [2]. If $F \in \mathbf{J}(X)$ also satisfies the condition $\inf_{x, y \in E} F(x, y) > -1$, then the process

$$t \mapsto \sum_{0 \leq s < t} F(X_{s-}, X_s) - \int_0^t \int_{\mathbb{R}^d} F(X_s, y) N(X_s, dy) dH_s$$

is a martingale and its Doleans-Dade exponential is (see Theorem 9.39 of [5])

$$\begin{aligned} L_t &= \exp \left(- \int_0^t \int_{\mathbb{R}^d} F(X_s, y) N(X_s, dy) dH_s \right) \prod_{s \leq t} (1 + F(X_{s-}, X_s)) \\ &= \exp \left(\sum_{s \leq t} \ln(1 + F(X_{s-}, X_s)) - \int_0^t \int_{\mathbb{R}^d} F(X_s, y) N(X_s, dy) dH_s \right). \end{aligned}$$

L_t is a nonnegative local martingale and thus a supermartingale multiplicative functional of X . L_t defines a family of probability of measures $\{\mathbb{P}_x, x \in \mathbb{R}^d\}$ on $\mathcal{M}_\infty$ by $d\tilde{\mathbb{P}}_x = L_t d\mathbb{P}_x$ on $\mathcal{M}_t$, see section 62 of [6]. Let $\tilde{X} = (\tilde{X}_t, \mathcal{M}, \mathcal{M}_t, \tilde{\mathbb{P}}_x, x \in \mathbb{R}^d)$ denote this new process. Under this family of measures $\tilde{\mathbb{P}}_x$, X_t is denoted by $\tilde{X}_t$. The process $\tilde{X}_t$ is called a purely discontinuous Girsanov transform of X . Under further assumption that $F \in \mathbf{A}_2(X)$ (see [3] for definition) and with a Lusin space E instead of $\mathbb{R}^d$, it was shown in [3] that the Green function of a purely discontinuous Girsanov transform of $\tilde{X}$ is comparable to that of X , especially when X is a symmetric stable process. Under the assumption that $F \in \mathbf{I}_2(X)$ (see [9] for definition) and with a Lusin space E instead of $\mathbb{R}^d$, it was shown in [9] that when X is a symmetric stable process, the transition density of a purely discontinuous Girsanov transform of $\tilde{X}$ is comparable to that of X .

One of the main tools used in [3] and [9] is the Dirichlet form. The Dirichlet form was also used in [8] to study the two-sided estimates on the density of the Feynman-Kac semigroups of symmetric α -stable-like processes (see definition 1.1). The Dirichlet form works well when the processes are symmetric. It encounters difficulty when the processes are not symmetric.

Let's introduce α -stable-like processes in the following.

Suppose the process $X = (X_t, \mathbb{P}_t)$ given above has a Lévy system (N, H) given by $H_t = t$ and

$$N(x, dy) = 2C(x, y)|x - y|^{-(d+\alpha)}m(dy),$$

where m a σ -finite measure on $\mathcal{B}(\mathbb{R}^d)$ given by $m(dx) = M(x)dx$ with $M(x)$ nonnegative and bounded.

Definition 1.1 *We say that X is an α -stable-like process if $C(x, y)$ is bounded.*

In this paper we assume that X admits a transition density $p(t, x, y)$ with respect to m and $p(t, x, y)$ is jointly continuously on $(0, \infty) \times \mathbb{R}^d \times \mathbb{R}^d$ and satisfies the condition

$$M_1 t^{-\frac{d}{\alpha}} \left(1 \wedge \frac{t^{\frac{1}{\alpha}}}{|x - y|}\right)^{d+\alpha} \leq p(t, x, y) \leq M_2 t^{-\frac{d}{\alpha}} \left(1 \wedge \frac{t^{\frac{1}{\alpha}}}{|x - y|}\right)^{d+\alpha}, \quad \forall (t, x, z) \in (0, \infty) \times \mathbb{R}^d \times \mathbb{R}^d, \quad (1.1)$$

where M_1 and M_2 are positive constants.

Here we do not assume that X is symmetric. When X is symmetric, it is called a symmetric α -stable-like process, which was introduced in [4], where a symmetric Hunt process is associated with a regular Dirichlet form and thus Dirichlet form method can be applied.

We list some examples for α -stable-like processes. One dimensional α -stable processes with their Lévy measure ν concentrated neither on $(0, \infty)$ nor on $(-\infty, 0)$ are α -stable-like processes, as their $C(x, y)$ are bounded and densities $p(t, x, y)$ satisfies (1.1), see Definition 14.16 and Remark 14.18

in [7]. For higher dimensions, [11] gave a large class of nonsymmetric strictly α -stable processes which satisfy (1.1) with $C(x, y)$ bounded. So this class also belongs to the class of α -stable-like processes.

In [10], we developed a method, which was based on an idea in [1] for Brownian motions, some results on discontinuous functionals and some techniques for integral estimates, to obtain the estimates on the density of the Feynman-Kac semigroups of α -stable-like processes where neither F nor X is symmetric. In this paper, by improving this method so that it also works for the purely discontinuous Girsanov transform, we will show that the transition density of the purely discontinuous Girsanov transform of $\tilde{X}$ is comparable to that of X with neither F nor X being symmetric.

The content of this paper is organized as follows. In section 2, we present preliminary results on additive functionals. In section 3, we establish the two-sided estimates on the transition density of the purely discontinuous Girsanov transformations under certain assumptions of $F(x, y)$.

2 Preliminary Results on Additive Functionals

For any $l > 0$, let $A_t^{(l)} = B_t^{(l)} - D_t^{(l)}$, where

$$B_t^{(l)} = \sum_{s \leq t} \ln(1 + F(X_{s-}, X_s)) 1_{\{|X_{s-} - X_s| > 1/l\}},$$

and

$$D_t^{(l)} = \int_0^t \int_E F(X_s, y) 1_{\{|X_s - y| > 1/l\}} N(X_s, dy) dH_s, \quad l = 1, 2, \dots$$

$B_t^{(l)}$ is a pure discontinuous functional and $D_t^{(l)}$ is a continuous functional. For convenience of notation, we omit the index l , i.e. denote $A_t^{(l)}$, $B_t^{(l)}$ and $D_t^{(l)}$ by A_t , B_t and D_t respectively. We have the following formulae for A_t^n .

Theorem 2.1

$$\begin{aligned} A_t^n &= C_n^1 \int_0^t A_s^{n-1} dA_s - C_n^2 \int_0^t A_s^{n-2} (\ln(1 + F(X_{s-}, X_s))) 1_{\{|X_{s-} - X_s| > 1/l\}} dA_s + \dots \\ &\quad + (-1)^{i-1} C_n^i \int_0^t A_s^{n-i} (\ln(1 + F(X_{s-}, X_s)))^{i-1} 1_{\{|X_{s-} - X_s| > 1/l\}} dA_s + \dots \\ &\quad + (-1)^{n-1} C_n^n \int_0^t (\ln(1 + F(X_{s-}, X_s)))^{n-1} 1_{\{|X_{s-} - X_s| > 1/l\}} dA_s \end{aligned}$$

and

$$\begin{aligned}
A_t^n &= C_n^1 \int_0^t (A_t - A_s)^{n-1} dA_s + C_n^2 \int_0^t (A_t - A_s)^{n-2} (\ln(1 + F(X_{s-}, X_s))) 1_{\{|X_{s-} - X_s| > 1/l\}} dA_s \\
&+ \cdots + C_n^i \int_0^t (A_t - A_s)^{n-i} (\ln(1 + F(X_{s-}, X_s)))^{i-1} 1_{\{|X_{s-} - X_s| > 1/l\}} dA_s + \cdots \\
&+ C_n^m \int_0^t (\ln(1 + F(X_{s-}, X_s)))^{n-1} 1_{\{|X_{s-} - X_s| > 1/l\}} dA_s.
\end{aligned}$$

Proof. Note that

$$A_s - A_{s-} = B_s - B_{s-} = \ln(1 + F(X_{s-}, X_s)) 1_{\{|X_{s-} - X_s| > 1/l\}}.$$

We use induction to show these two formulae for A_t^n . It is clear that they hold for $n = 2$. Suppose they hold for $n \leq m - 1$, we show they hold for $n = m$.

It follows from the integration by parts formula,

$$A_t^m = \int_0^t A_{s-} dA_s^{m-1} + \int_0^t A_s^{m-1} dA_s,$$

where

$$\begin{aligned}
&\int_0^t A_{s-} dA_s^{m-1} \\
&= \int_0^t (A_s - \ln(1 + F(X_{s-}, X_s)) 1_{\{|X_{s-} - X_s| > 1/l\}}) dA_s^{m-1} \\
&= \int_0^t A_s dA_s^{m-1} - \int_0^t \ln(1 + F(X_{s-}, X_s)) 1_{\{|X_{s-} - X_s| > 1/l\}} dA_s^{m-1} \\
&= \int_0^t A_s \left(\sum_{i=1}^{m-1} (-1)^{i-1} C_{m-1}^i A_s^{m-1-i} (\ln(1 + F(X_{s-}, X_s)) 1_{\{|X_{s-} - X_s| > 1/l\}})^{i-1} \right) dA_s \\
&\quad - \int_0^t F(X_{s-}, X_s) \left(\sum_{j=1}^{m-1} (-1)^{j-1} C_{m-1}^j A_s^{m-1-j} (\ln(1 + F(X_{s-}, X_s)) 1_{\{|X_{s-} - X_s| > 1/l\}})^{j-1} \right) dA_s \\
&\quad \cdot dA_s \text{ (by the first formula for } A_t^n \text{ when } n = m - 1 \text{)} \\
&= \sum_{i=1}^{m-1} (-1)^{i-1} C_{m-1}^i \int_0^t A_s^{m-i} (\ln(1 + F(X_{s-}, X_s)) 1_{\{|X_{s-} - X_s| > 1/l\}})^{i-1} dA_s \\
&\quad - \sum_{j=1}^{m-1} (-1)^{j-1} C_{m-1}^j \int_0^t A_s^{m-1-j} (\ln(1 + F(X_{s-}, X_s)) 1_{\{|X_{s-} - X_s| > 1/l\}})^{j-1} dA_s
\end{aligned}$$

$$\begin{aligned}
&= \sum_{i=1}^{m-1} (-1)^{i-1} C_{m-1}^i \int_0^t A_s^{m-i} (\ln(1 + F(X_{s-}, X_s)) 1_{\{|X_{s-} - X_s| > 1/l\}})^{i-1} dA_s \\
&\quad - \sum_{i=2}^m (-1)^{i-2} C_{m-1}^{i-1} \int_0^t A_s^{m-i} (\ln(1 + F(X_{s-}, X_s)) 1_{\{|X_{s-} - X_s| > 1/l\}})^{i-1} dA_s \\
&\quad (\text{ let } j = i - 1) \\
&= \sum_{i=2}^{m-1} (-1)^{i-1} (C_{m-1}^i + C_{m-1}^{i-1}) \int_0^t A_s^{m-i} (\ln(1 + F(X_{s-}, X_s)) 1_{\{|X_{s-} - X_s| > 1/l\}})^{i-1} dA_s \\
&\quad + C_{m-1}^1 \int_0^t A_s^{m-1} dA_s - (-1)^{m-2} \int_0^t (\ln(1 + F(X_{s-}, X_s)) 1_{\{|X_{s-} - X_s| > 1/l\}})^{m-1} dA_s \\
&= \sum_{i=2}^{m-1} (-1)^{i-1} C_m^i \int_0^t A_s^{m-i} (\ln(1 + F(X_{s-}, X_s)) 1_{\{|X_{s-} - X_s| > 1/l\}})^{i-1} dA_s \\
&\quad + C_{m-1}^1 \int_0^t A_s^{m-1} dA_s - (-1)^m \int_0^t (\ln(1 + F(X_{s-}, X_s)) 1_{\{|X_{s-} - X_s| > 1/l\}})^{m-1} dA_s \\
&\quad (\text{ by } C_{m-1}^i + C_{m-1}^{i-1} = C_m^i).
\end{aligned}$$

Thus

$$\begin{aligned}
A_t^m &= \int_0^t A_{s-} dA_s^{m-1} + \int_0^t A_s^{m-1} dA_s \\
&= \sum_{i=1}^m (-1)^{i-1} C_m^i \int_0^t A_s^{m-i} (\ln(1 + F(X_{s-}, X_s)) 1_{\{|X_{s-} - X_s| > 1/l\}})^{i-1} dA_s,
\end{aligned}$$

i.e. the first formula for A_t^n holds for $n=m$.

Now we go to the second formula for A_t^n , for $n = m$.

$$\begin{aligned}
&C_m^1 \int_0^t (A_t - A_s)^{m-1} dA_s \\
&= C_m^1 \int_0^t \sum_{i=0}^{m-1} C_{m-1}^i A_t^i (-1)^{m-1-i} A_s^{m-1-i} dA_s \\
&= \sum_{i=0}^{m-1} (-1)^{m-1-i} C_m^1 C_{m-1}^i A_t^i \int_0^t A_s^{m-1-i} dA_s \\
&= \sum_{i=0}^{m-1} (-1)^{m-1-i} C_m^i (m-i) A_t^i \int_0^t A_s^{m-1-i} dA_s \\
&\quad (\text{ by } C_m^1 C_{m-1}^i = C_m^i (m-i)) \\
&= \sum_{i=0}^{m-1} (-1)^{m-1-i} C_m^i A_t^i ((m-i) \int_0^t A_s^{m-1-i} dA_s)
\end{aligned}$$

$$\begin{aligned}
&= \sum_{i=0}^{m-1} (-1)^{m-1-i} C_m^i A_t^i (A_t^{m-i} + \int_0^t \sum_{k=2}^{m-i} (-1)^k C_{m-i}^k A_s^{m-i-k} (\ln(1 + F(X_{s-}, X_s)) \\
&\quad \cdot 1_{\{|X_{s-} - X_s| > 1/l\}})^{k-1} dA_s) \\
&\quad (\text{ by the first formula of } A_t^n \text{ for } n = m - i) \\
&= \sum_{i=0}^{m-1} (-1)^{m-1-i} C_m^i A_t^i A_t^{m-i} + \int_0^t \sum_{i=0}^{m-1} (-1)^{m-1-i} \sum_{k=2}^{m-i} (-1)^k C_m^i C_{m-i}^k A_t^i A_s^{m-i-k} \\
&\quad \cdot (\ln(1 + F(X_{s-}, X_s)) 1_{\{|X_{s-} - X_s| > 1/l\}})^{k-1} dA_s,
\end{aligned}$$

where

$$\sum_{i=0}^{m-1} (-1)^{m-1-i} C_m^i A_t^i A_t^{m-i} = \left(\sum_{i=0}^{m-1} (-1)^{m-1-i} C_m^i \right) A_t^m = (1) A_t^m,$$

and

$$\begin{aligned}
&\int_0^t \sum_{i=0}^{m-1} (-1)^{m-1-i} \sum_{k=2}^{m-i} (-1)^k C_m^i C_{m-i}^k A_t^i A_s^{m-i-k} (\ln(1 + F(X_{s-}, X_s)) 1_{\{|X_{s-} - X_s| > 1/l\}})^{k-1} dA_s \\
&= \int_0^t \sum_{k=2}^m \sum_{i=0}^{m-k} (-1)^{m-k-i-1} C_m^k C_{m-k}^i A_t^i A_s^{m-k-i} (\ln(1 + F(X_{s-}, X_s)) 1_{\{|X_{s-} - X_s| > 1/l\}})^{k-1} dA_s \\
&\quad (\text{ by } C_m^i C_{m-i}^k = C_m^k C_{m-k}^i \text{ and } (-1)^{m-1-i+k} = (-1)^{m-k-i-1}) \\
&= \sum_{k=2}^m C_m^k (-1)^{-1} \int_0^t (A_t - A_s)^{m-k} (\ln(1 + F(X_{s-}, X_s)) 1_{\{|X_{s-} - X_s| > 1/l\}})^{k-1} dA_s \\
&= - \sum_{k=2}^m C_m^k \int_0^t (A_t - A_s)^{m-k} (\ln(1 + F(X_{s-}, X_s)) 1_{\{|X_{s-} - X_s| > 1/l\}})^{k-1} dA_s,
\end{aligned}$$

therefore

$$\begin{aligned}
&C_m^1 \int_0^t (A_t - A_s)^{m-1} dA_s \\
&= A_t^m - \sum_{k=2}^m C_m^k \int_0^t (A_t - A_s)^{m-k} (\ln(1 + F(X_{s-}, X_s)) 1_{\{|X_{s-} - X_s| > 1/l\}})^{k-1} dA_s,
\end{aligned}$$

i.e. the second formula for A_t^n holds for $n = m$. □

3 Transition Density of Pure Jump Girsanov Transformation

We need the following lemmas to prove the main results.

Lemma 3.1 For any two positive constants $K < 1$ and L , there exist two constants $C_{01}(K, L)$ and $C_{02}(K, L)$ which depend on K and L , such that

$$\begin{aligned} L^{-1}K^{n-1} + \frac{K^{n-2}}{2!} + \frac{LK^{n-3}}{3!} + \cdots + \frac{L^{i-2}K^{n-i}}{i!} + \cdots + \frac{L^{n-2}}{n!} &\leq C_{01}(K, L)K^n, \\ LK^{n-1} + \frac{L^2K^{n-2}}{2!} + \frac{L^3K^{n-3}}{3!} + \cdots + \frac{L^iK^{n-i}}{i!} + \cdots + \frac{L^n}{n!} &\leq C_{02}(K, L)K^n, \text{ for any } n \geq 1. \end{aligned}$$

Proof. Use the fact that there exists $i_0 \geq 0$, such that when $j \geq i_0$,

$$\frac{L^j}{j!} \leq \left(\frac{K}{2}\right)^j.$$

□

Lemma 3.2 For any two positive constants $K < 1$ and L , there exists a constant $C_{03}(K, L)$ which depends on K and L , such that

$$\sum_{i=2}^m \sum_{j=2}^{m-i} C_m^i C_{m-i}^j (m-i-j)! L^{i-2} L^{j-2} K^{m-i-j} \leq C_{03}(K, L) m! K^m, \text{ for all } m \geq 2.$$

Proof.

$$\begin{aligned} &\sum_{i=2}^m \sum_{j=2}^{m-i} C_m^i C_{m-i}^j (m-i-j)! L^{i-2} L^{j-2} K^{m-i-j} \\ &= \sum_{i=2}^m \sum_{j=2}^{m-i} \frac{m!}{i!j!} L^{i-2} L^{j-2} K^{m-i-j} \\ &= m! \sum_{i=2}^m \frac{L^{i-2}}{i!} \left(\sum_{j=2}^{m-i} \frac{L^{j-2}}{j!} K^{m-i-j} \right) \\ &\leq m! \sum_{i=2}^m \frac{L^{i-2}}{i!} \tilde{C}_{01}(L, K) K^{m-i} \\ &\quad \text{(by the same argument of lemma 3.1, where } \tilde{C}_{01}(L, K) \text{ is a positive constant)} \\ &\leq C_{03}(L, K) m! K^m \\ &\quad \text{(by the same argument of lemma 3.1, where } \tilde{C}_{03}(L, K) \text{ is a positive constant)}. \end{aligned}$$

□

From now on we define $q_0(t, x, y) = p(t, x, y)$ where $p(t, x, y)$ is the transition density of α -stable-like process X and satisfies (1.1).

By the second formula for A_t^n , we have for any g bounded measurable

$$\begin{aligned}
& \mathbb{E}_x[A_t^n g(X_t)] \\
&= \sum_{i=1}^n C_n^i \mathbb{E}_x \left[\int_0^t (A_t - A_s)^{n-i} g(X_t) (\ln(1 + F(X_{s-}, X_s)))^{i-1} 1_{\{|X_{s-} - X_s| > 1/l\}} dA_s \right] \\
&= \sum_{i=1}^n C_n^i \mathbb{E}_x \left[\int_0^t (A_t - A_s)^{n-i} g(X_t) (\ln(1 + F(X_{s-}, X_s)))^{i-1} 1_{\{|X_{s-} - X_s| > 1/l\}} dB_s \right] \\
&\quad - \sum_{i=1}^n C_n^i \mathbb{E}_x \left[\int_0^t (A_t - A_s)^{n-i} g(X_t) (\ln(1 + F(X_{s-}, X_s)))^{i-1} 1_{\{|X_{s-} - X_s| > 1/l\}} dD_s \right] \\
&= \sum_{i=1}^n C_n^i \mathbb{E}_x \left[\int_0^t \mathbb{E}_{X_s} (A_{t-s}^{n-i} g(X_{t-s})) d \left(\sum_{r \leq s} (\ln(1 + F(X_{r-}, X_r)))^i 1_{\{|X_{r-} - X_r| > 1/l\}} \right) \right] \\
&\quad - C_n^1 \mathbb{E}_x \left[\int_0^t \mathbb{E}_{X_s} (A_{t-s}^{n-1} g(X_{t-s})) dD_s \right] \\
&= \sum_{i=1}^n C_n^i \mathbb{E}_x \left[\int_0^t \int_{\mathbb{R}^d} \frac{2C(X_s, y) (\ln(1 + F(X_s, y)))^{i-1} 1_{\{|X_s - y| > 1/l\}}}{|X_s - y|^{d+\alpha}} \mathbb{E}_y (A_{t-s}^{n-i} g(X_{t-s})) m(dy) ds \right] \\
&\quad - C_n^1 \mathbb{E}_x \left[\int_0^t \int_{\mathbb{R}^d} \frac{2C(X_s, y) F(X_s, y) 1_{\{|X_s - y| > 1/l\}}}{|X_s - y|^{d+\alpha}} \mathbb{E}_y (A_{t-s}^{n-1} g(X_{t-s})) m(dy) ds \right] \\
&= C_n^1 \mathbb{E}_x \left[\int_0^t \int_{\mathbb{R}^d} \frac{2C(X_s, y) (\ln(1 + F(X_s, y))) - F(X_s, y) 1_{\{|X_s - y| > 1/l\}}}{|X_s - y|^{d+\alpha}} \mathbb{E}_y (A_{t-s}^{n-1} g(X_{t-s})) m(dy) \right. \\
&\quad \cdot ds \left. + \sum_{i=2}^n C_n^i \mathbb{E}_x \left[\int_0^t \int_{\mathbb{R}^d} \frac{2C(X_s, y) (\ln(1 + F(X_s, y)))^{i-1} 1_{\{|X_s - y| > 1/l\}}}{|X_s - y|^{d+\alpha}} \mathbb{E}_y (A_{t-s}^{n-i} g(X_{t-s})) m(dy) \right. \right. \\
&\quad \cdot ds \left. \right].
\end{aligned}$$

We define $q_n^{(l)}(t, x, z)$ as follows,

$$\begin{aligned}
& q_n^{(l)}(t, x, z) \\
&= C_n^1 \int_0^t \int_{\mathbb{R}^d} p(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{2C(w, y) (\ln(1 + F(w, y))) - F(w, y) 1_{\{|w - y| > 1/l\}}}{|w - y|^{d+\alpha}} \\
&\quad \cdot q_{n-1}(t - s, y, z) m(dy) ds + \sum_{i=2}^n C_n^i \int_0^t \int_{\mathbb{R}^d} p(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{2C(w, y) (\ln(1 + F(w, y)))^i}{|w - y|^{d+\alpha}} \\
&\quad \cdot 1_{\{|w - y| > 1/l\}} q_{n-i}(t - s, y, z) m(dy) ds.
\end{aligned}$$

For convenience, we denote $q_n^{(l)}(t, x, z)$ by $q_n(t, x, z)$.

Then by induction, we can show that for any $n \geq 1$,

$$\int_{\mathbb{R}^d} q_n(t, x, z) g(z) m(dz) = \mathbb{E}_x[A_t^n g(X_t)],$$

and

$$\begin{aligned} & \mathbb{E}_x[A_t^n g(X_t)] \\ &= C_n^1 \mathbb{E}_x \left[\int_0^t \int_{\mathbb{R}^d} \frac{2C(X_s, y)(\ln(1 + F(X_s, y)) - F(X_s, y)) 1_{\{|X_s - y| > 1/l\}}}{|X_s - y|^{d+\alpha}} \int_{\mathbb{R}^d} q_{n-1}(t-s, y, z) g(z) \right. \\ & \quad \cdot m(dz) m(dy) ds \Big] + \sum_{i=2}^n C_n^i \mathbb{E}_x \left[\int_0^t \int_{\mathbb{R}^d} \frac{2C(X_s, y)(\ln(1 + F(X_s, y)))^i 1_{\{|X_s - y| > 1/l\}}}{|X_s - y|^{d+\alpha}} \right. \\ & \quad \cdot \int_{\mathbb{R}^d} q_{n-i}(t-s, y, z) g(z) m(dz) m(dy) ds \Big] \\ &= C_n^1 \int_0^t \int_{\mathbb{R}^d} p(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{2C(w, y)(\ln(1 + F(w, y)) - F(w, y)) 1_{\{|w - y| > 1/l\}}}{|w - y|^{d+\alpha}} \\ & \quad \cdot \int_{\mathbb{R}^d} q_{n-1}(t-s, y, z) g(z) m(dz) m(dy) ds + \sum_{i=2}^n C_n^i \int_0^t \int_{\mathbb{R}^d} p(s, x, w) m(dw) \\ & \quad \cdot \int_{\mathbb{R}^d} \frac{2C(w, y)(\ln(1 + F(w, y)))^i 1_{\{|w - y| > 1/l\}}}{|w - y|^{d+\alpha}} \cdot \int_{\mathbb{R}^d} q_{n-i}(t-s, y, z) g(z) m(dz) m(dy) ds. \end{aligned}$$

Let $\mathbf{K}_{\mathbf{d}, \alpha}$ be the Kato class. Recall the definition of Kato class. We say that a function V on $\mathbb{R}^d$ belongs to the Kato class $\mathbf{K}_{\mathbf{d}, \alpha}$ if $\lim_{t \downarrow 0} \sup_{x \in \mathbb{R}^d} \int_0^t \int_{\mathbb{R}^d} p(t, x, y) |V(y)| dy ds = 0$. We say that a signed measure μ on $\mathbb{R}^d$ belongs to the Kato class $\mathbf{K}_{\mathbf{d}, \alpha}$ if $\lim_{t \downarrow 0} \sup_{x \in \mathbb{R}^d} \int_0^t \int_{\mathbb{R}^d} p(t, x, y) |\mu|(dy) ds = 0$.

Suppose F is a function on $\mathbb{R}^d \times \mathbb{R}^d$.

Definition 3.1 We say F belongs to $\mathbf{J}_{\mathbf{d}, \alpha}$ if F is bounded, vanishing on the diagonal, and the function

$$x \mapsto \int_{\mathbb{R}^d} \frac{|F(x, y)|}{|x - y|^{d+\alpha}} dy$$

belongs to $\mathbf{K}_{\mathbf{d}, \alpha}$.

We assume $\inf_{x, y \in E} F(x, y) > -1$, $F(x, y)$ vanishes on the diagonal and satisfies

$$w \mapsto \int_{\mathbb{R}^d} \frac{F^2(w, y)}{|w - y|^{d+\alpha}} dy$$

and

$$y \mapsto \int_{\mathbb{R}^d} \frac{F^2(w, y)}{|w - y|^{d+\alpha}} dw$$

both belong to $\mathbf{K}_{\mathbf{d}, \alpha}$, i.e. $F^2(x, y)$ and $F^2(y, x)$ both belongs to $\mathbf{J}_{\mathbf{d}, \alpha}$.

It is clear that there exist constants C_1 and C_2 such that

$$|\ln(1 + F(x, y)) - F(x, y)| \leq C_1 F^2(x, y), \quad \forall x, y \in E,$$

and

$$|\ln(1 + F(x, y))| \leq C_2 |F(x, y)|, \quad \forall x, y \in E.$$

For the constants C_1 and C_2 given above, we assume that there exist constants $\overline{C}$, L and $\overline{M}$ such that

$$\max(|2C(x, y)C_1|, |2C(x, y)C_2|, C_2) \leq \overline{C} \quad (3.1)$$

$$\max(|2F(x, y)|, |2\overline{C}F(x, y)|) \leq L, \quad (3.2)$$

and $0 < M(y) \leq \overline{M}$ where $m(dy) = M(y)dy$.

Define $\overline{F}(x, y) = |F(x, y)| + |F(y, x)|$, which is symmetric and satisfies $|\overline{F}(w, y)| \leq L$. Define $\overline{p}(t, x, y) = p(t, x, y) + p(t, y, x)$. Then $\overline{p}(t, x, y)$ is symmetric and satisfies

$$2M_1 t^{-\frac{d}{\alpha}} \left(1 \wedge \frac{t^{\frac{1}{\alpha}}}{|x - y|}\right)^{d+\alpha} \leq \overline{p}(t, x, y) \leq 2M_2 t^{-\frac{d}{\alpha}} \left(1 \wedge \frac{t^{\frac{1}{\alpha}}}{|x - y|}\right)^{d+\alpha}, \quad \forall (t, x, y) \in (0, \infty) \times \mathbb{R}^d \times \mathbb{R}^d.$$

Denote $(\int_{\mathbb{R}^d} \frac{\overline{F}^2(w, y)}{|w - y|^{d+\alpha}} dy)dw$ by $\mu(dw)$ and let $C_t = \sup_{x \in \mathbb{R}^d} \int_0^t \int_{\mathbb{R}^d} \overline{p}(s, x, w) \mu(dw) ds$. Then $C_t \downarrow 0$ as $t \downarrow 0$. It is clear that there exist two positive constants D_1 and D_2 such that $D_1 \leq \int_{\mathbb{R}^d} \overline{p}(t, x, y) m(dy) \leq D_2$, as $\overline{p}(t, x, y)$ is comparable to $p(t, x, y)$. Let $\overline{q}_0(t, x, z) = \overline{p}(s, x, z)$ and define $\overline{q}_n(t, x, z)$ by

$$\begin{aligned} \overline{q}_n(t, x, z) &= C_n^1 \int_0^t \int_{\mathbb{R}^d} \overline{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{C} \overline{F}^2(w, y)}{|w - y|^{d+\alpha}} \overline{q}_{n-1}(t - s, y, z) m(dy) ds \\ &\quad + \sum_{i=2}^n C_n^i \int_0^t \int_{\mathbb{R}^d} \overline{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{C}^i \overline{F}^i(w, y)}{|w - y|^{d+\alpha}} \overline{q}_{n-i}(t - s, y, z) m(dy) ds. \end{aligned}$$

By the choice of $\overline{C}$ in (3.1), it is clear that $|q_n(t, x, z)| \leq \overline{q}_n(t, x, z)$. We can see that $q_n(t, x, z)$ may not be symmetric even when $F(x, z)$ and $p(t, x, z)$ are symmetric. Thus we can not use an argument similar as that used in lemma 3.3 in [10].

Next are the main results.

For the positive constants L , $\overline{C}$, and $\overline{M}$ given above, and $K < 1$, M_2 given in Lemma 3.1, Lemma 3.2 and (1.1), there exists positive constants $\tilde{C}$ and $\tilde{C}_1$ such that

$$\begin{aligned}\overline{C}\overline{M}^2 D_2 &\leq \tilde{C}K, \\ L^{m-2}\overline{C}^2\overline{M}^2 &\leq \frac{1}{2}\tilde{C}m!K^m, \quad \forall m > 0, \\ \overline{C}\frac{L^2 D_2^2}{(1 - \frac{1}{10} - 2^{-\frac{1}{2}})^{d+\alpha}} &\leq \frac{1}{4}\tilde{C}_1 K, \quad \forall m > 0 \\ \overline{C}\frac{L^m D_2^2}{(1 - \frac{1}{10} - 2^{-\frac{1}{2}})^{d+\alpha}} &\leq \frac{1}{4}\tilde{C}_1 m!K^m, \quad \forall m > 0.\end{aligned}$$

We claim

Theorem 3.3 *There exist constants $t_2 > 0$, $\tilde{C} > 0$ and $\tilde{C}_1 > 0$ such that when $0 < t \leq t_2$,*

$$1. \quad \int_{\mathbb{R}^d} \overline{q}_n(t, x, z) m(dz) \leq \tilde{C} C_t n! K^n, \quad \forall n \geq 1, \quad (3.3)$$

$$\int_{\mathbb{R}^d} \overline{q}_n(t, x, z) m(dx) \leq \tilde{C} C_t n! K^n, \quad \forall n \geq 1. \quad (3.4)$$

$$2. \quad \overline{q}_n(t, x, z) \leq \tilde{C}_1 n! K^n t^{-\frac{d}{\alpha}} \left(1 \wedge \frac{t^{\frac{1}{\alpha}}}{|x - z|} \right)^{d+\alpha}, \quad \forall n \geq 0 \quad (3.5)$$

Proof. We show statement 1 holds for $n = 1$

$$\begin{aligned}&\int_{\mathbb{R}^d} \overline{q}_1(t, x, z) m(dz) \\&\leq \int_{\mathbb{R}^d} \int_0^t \int_{\mathbb{R}^d} \overline{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{CF}^2(w, y)}{|w - y|^{d+\alpha}} \overline{p}(t - s, y, z) m(dy) ds m(dz) \\&= \int_0^t \int_{\mathbb{R}^d} \overline{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{CF}^2(w, y)}{|w - y|^{d+\alpha}} \int_{\mathbb{R}^d} \overline{p}(t - s, y, z) m(dz) m(dy) ds \\&= \int_0^t \int_{\mathbb{R}^d} \overline{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{CF}^2(w, y)}{|w - y|^{d+\alpha}} m(dy) ds D_2 \\&\leq \overline{C}\overline{M}^2 C_t D_2 \leq \tilde{C} C_t K\end{aligned}$$

and

$$\begin{aligned}&\int_{\mathbb{R}^d} \overline{q}_1(t, x, z) m(dx) \\&\leq \int_{\mathbb{R}^d} \int_0^t \int_{\mathbb{R}^d} \overline{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{CF}^2(w, y)}{|w - y|^{d+\alpha}} \overline{p}(t - s, y, z) m(dy) ds m(dx)\end{aligned}$$

$$\begin{aligned}
&= \int_0^t \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \bar{p}(s, x, w) m(dx) m(dw) \int_{\mathbb{R}^d} \frac{\overline{CF}^2(w, y)}{|w - y|^{d+\alpha}} \bar{p}(t - s, y, z) m(dy) ds \\
&= \int_0^t \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \bar{p}(s, w, x) m(dx) m(dw) \int_{\mathbb{R}^d} \frac{\overline{CF}^2(w, y)}{|w - y|^{d+\alpha}} \bar{p}(t - s, y, z) m(dy) ds \\
&\quad (\text{by symmetry of } \bar{p}(s, x, w)) \\
&= \int_0^t \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \frac{\overline{CF}^2(w, y)}{|w - y|^{d+\alpha}} m(dw) p(t - s, y, z) m(dy) ds \\
&\leq \overline{CM}^2 C_t D_2 \leq \tilde{C} C_t K.
\end{aligned}$$

Thus statement 1 holds for $n = 1$.

Since $\bar{q}_0(t, x, z) = \bar{p}(t, x, z)$, statement 2 holds for $n = 0$. We show it also holds for $n = 1$. We write $\bar{q}_1(t, x, z)$ in the following way,

$$\begin{aligned}
\bar{q}_1(t, x, z) &= \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \bar{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{CF}^2(w, y)}{|w - y|^{d+\alpha}} \bar{p}(t - s, y, z) m(dy) ds \\
&\quad + \int_{\frac{t}{2}}^t \int_{\mathbb{R}^d} \bar{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{CF}^2(w, y)}{|w - y|^{d+\alpha}} \bar{p}(t - s, y, z) m(dy) ds.
\end{aligned}$$

First we look at the first term in the expression of $\bar{q}_1(t, x, z)$.

There are two cases:

Case 1. When $|x - z| \leq t^{\frac{1}{\alpha}}$,

$$\begin{aligned}
&\int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \bar{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{CF}^2(w, y)}{|w - y|^{d+\alpha}} \bar{p}(t - s, y, z) m(dy) ds \\
&\leq \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \bar{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{F}^2(w, y)}{|w - y|^{d+\alpha}} dy ds \overline{CM} \tilde{C}_1 \left(\frac{t}{2}\right)^{-\frac{d}{\alpha}} \\
&\leq C_t \overline{CM}^2 \tilde{C}_1 \left(\frac{t}{2}\right)^{-\frac{d}{\alpha}}.
\end{aligned}$$

Since $C_t \downarrow 0$ as $t \downarrow 0$, there exists $t_{01} > 0$ such that when $0 < t \leq t_{01}$,

$$C_t \overline{CM}^2 \left(\frac{1}{2}\right)^{-\frac{d}{\alpha}} \leq \frac{1}{2} K.$$

Thus

$$\int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \bar{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{CF}^2(w, y)}{|w - y|^{d+\alpha}} \bar{p}(t - s, y, z) m(dy) ds \leq \frac{1}{2} \tilde{C}_1 K t^{-\frac{d}{\alpha}}.$$

Case 2. When $|x - z| \geq t^{\frac{1}{\alpha}}$. Let $B_1 = \{y \in \mathbb{R}^d \mid |y - z| \geq \frac{1}{10}|x - z|\}$, $B_2 = \{w \in \mathbb{R}^d \mid |w - x| \geq 2^{-\frac{1}{2}}|x - z|\}$ and $B_3 = \{(w, y) \in \mathbb{R}^d \times \mathbb{R}^d \mid |y - z| < \frac{1}{10}|x - z|, |w - x| < 2^{-\frac{1}{2}}|x - z|\}$. On B_3 , we have $|w - y| \geq (1 - \frac{1}{10} - 2^{-\frac{1}{2}})|x - z|$.

$$\begin{aligned}
& \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \bar{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{CF}^2(w, y)}{|w - y|^{d+\alpha}} \bar{p}(t - s, y, z) m(dy) ds \\
& \leq \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \bar{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{CF}^2(w, y)}{|w - y|^{d+\alpha}} \bar{p}(t - s, y, z) 1_{B_1}(y) m(dy) ds \\
& \quad + \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \bar{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{CF}^2(w, y)}{|w - y|^{d+\alpha}} \bar{p}(t - s, y, z) 1_{B_2}(w) m(dy) ds \\
& \quad + \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \bar{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{CF}^2(w, y)}{|w - y|^{d+\alpha}} \bar{p}(t - s, y, z) 1_{B_3}(w, y) m(dy) ds \\
& \leq \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \bar{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{F}^2(w, y)}{|w - y|^{d+\alpha}} \overline{MC} \tilde{C}_1 10^{d+\alpha} \frac{(t - s)}{|x - z|^{d+\alpha}} dy ds \\
& \quad + \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} dw \tilde{C}_1 \overline{M}^2 2^{\frac{1}{2}(d+\alpha)} \frac{s}{|x - z|^{d+\alpha}} \int_{\mathbb{R}^d} \frac{\overline{F}^2(w, y)}{|w - y|^{d+\alpha}} \overline{C} \bar{p}(t - s, y, z) dy ds \\
& \quad + \overline{C} \frac{L^2}{(1 - \frac{1}{10} - 2^{-\frac{1}{2}})^{d+\alpha}} \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \bar{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{1}{|x - z|^{d+\alpha}} \bar{p}(t - s, y, z) m(dy) ds \\
& \leq \left(\tilde{C}_1 (\overline{CM}^2 10^{d+\alpha}) C_t + \tilde{C}_1 (\overline{CM}^2 2^{\frac{1}{2}(d+\alpha)}) C_t \right) \frac{t}{|x - z|^{d+\alpha}} + \overline{C} \frac{L^2 D_2^2}{(1 - \frac{1}{10} - 2^{-\frac{1}{2}})^{d+\alpha}} \frac{t}{|x - z|^{d+\alpha}} \\
& \quad (\text{by symmetry, } \bar{p}(t - s, y, z) = \bar{p}(t - s, z, y)).
\end{aligned}$$

There exists $t_{02} > 0$ such that when $0 < t \leq t_{02}$, C_t is small enough and the sum of the first two terms $\leq \frac{1}{4} \tilde{C}_1 K \frac{t}{|x - z|^{d+\alpha}}$. For the third term, we know that

$$\overline{C} \frac{L^2 D_2^2}{(1 - \frac{1}{10} - 2^{-\frac{1}{2}})^{d+\alpha}} \leq \frac{1}{4} \tilde{C}_1 K.$$

Thus

$$\int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \bar{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{CF}^2(w, y)}{|w - y|^{d+\alpha}} \bar{p}(t - s, y, z) m(dy) ds \leq \frac{1}{2} \tilde{C}_1 K \frac{t}{|x - z|^{d+\alpha}}.$$

Combining case 1 and case 2, when $0 < t \leq \min(t_{01}, t_{02})$,

$$\int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \bar{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{CF}^2(w, y)}{|w - y|^{d+\alpha}} \bar{p}(t - s, y, z) m(dy) ds \leq \frac{1}{2} \tilde{C}_1 K t^{-\frac{d}{\alpha}} \left(1 \wedge \frac{t^{\frac{1}{\alpha}}}{|x - z|} \right)^{d+\alpha}.$$

For the second term in the expression of $\bar{q}_1(t, x, z)$:

$$\int_{\frac{t}{2}}^t \int_{\mathbb{R}^d} \bar{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{CF}^2(w, y)}{|w - y|^{d+\alpha}} \bar{p}(t - s, y, z) m(dy) ds,$$

let $t - s = \tilde{s}$. The second term becomes

$$\int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \bar{p}(t - \tilde{s}, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{CF}^2(w, y)}{|w - y|^{d+\alpha}} \bar{p}(\tilde{s}, y, z) m(dy) d\tilde{s}.$$

There are two cases:

Case a. When $|x - z| \leq t^{\frac{1}{\alpha}}$,

$$\begin{aligned} & \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \bar{p}(t - \tilde{s}, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{CF}^2(w, y)}{|w - y|^{d+\alpha}} \bar{p}(\tilde{s}, y, z) m(dy) d\tilde{s} \\ & \leq \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \tilde{C}_1 \left(\frac{t}{2}\right)^{-\frac{d}{\alpha}} m(dw) \int_{\mathbb{R}^d} \frac{\overline{F}^2(w, y)}{|w - y|^{d+\alpha}} \bar{C} \bar{p}(\tilde{s}, z, y) m(dy) d\tilde{s} \\ & \leq \tilde{C}_1 \overline{CM}^2 \left(\frac{t}{2}\right)^{-\frac{d}{\alpha}} C_t. \end{aligned}$$

Since $C_t \downarrow 0$ as $t \downarrow 0$, there exists $t_{03} > 0$ such that when $0 < t \leq t_{03}$,

$$\overline{CM}^2 \left(\frac{1}{2}\right)^{-\frac{d}{\alpha}} C_t \leq \frac{1}{2} K.$$

Thus

$$\int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \bar{p}(t - \tilde{s}, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{CF}^2(w, y)}{|w - y|^{d+\alpha}} \bar{p}(\tilde{s}, y, z) m(dy) d\tilde{s} \leq \frac{1}{2} \tilde{C}_1 K t^{-\frac{d}{\alpha}}.$$

Case b. When $|x - z| \geq t^{\frac{1}{\alpha}}$. Let $B_1 = \{y \in \mathbb{R}^d \mid |y - z| \geq \frac{1}{10}|x - z|\}$, $B_2 = \{w \in \mathbb{R}^d \mid |w - x| \geq 2^{-\frac{1}{2}}|x - z|\}$ and $B_3 = \{(w, y) \in \mathbb{R}^d \times \mathbb{R}^d \mid |y - z| < \frac{1}{10}|x - z|, |w - x| < 2^{-\frac{1}{2}}|x - z|\}$. On B_3 , we have $|w - y| \geq (1 - \frac{1}{10} - 2^{-\frac{1}{2}})|x - z|$.

$$\begin{aligned} & \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \bar{p}(t - \tilde{s}, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{CF}^2(w, y)}{|w - y|^{d+\alpha}} \bar{p}(\tilde{s}, y, z) m(dy) d\tilde{s} \\ & \leq \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \bar{p}(t - \tilde{s}, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{CF}^2(w, y)}{|w - y|^{d+\alpha}} \bar{p}(\tilde{s}, y, z) 1_{B_1}(y) m(dy) d\tilde{s} \\ & \quad + \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \bar{p}(t - \tilde{s}, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{CF}^2(w, y)}{|w - y|^{d+\alpha}} \bar{p}(\tilde{s}, y, z) 1_{B_2}(w) m(dy) d\tilde{s} \\ & \quad + \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \bar{p}(t - \tilde{s}, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{CF}^2(w, y)}{|w - y|^{d+\alpha}} \bar{p}(\tilde{s}, y, z) 1_{B_3}(w, y) m(dy) d\tilde{s} \\ & \leq \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \bar{p}(t - \tilde{s}, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{F}^2(w, y)}{|w - y|^{d+\alpha}} \bar{C} \tilde{C}_1 10^{d+\alpha} \frac{\tilde{s}}{|x - z|^{d+\alpha}} m(dy) d\tilde{s} \\ & \quad + \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} dw \tilde{C}_1 \overline{M}^2 2^{\frac{1}{2}(d+\alpha)} \frac{t - \tilde{s}}{|x - z|^{d+\alpha}} \int_{\mathbb{R}^d} \frac{\overline{F}^2(w, y)}{|w - y|^{d+\alpha}} \bar{C} \bar{p}(\tilde{s}, z, y) dy d\tilde{s} \end{aligned}$$

$$\begin{aligned}
& + \overline{C} \frac{L^2}{(1 - \frac{1}{10} - 2^{-\frac{1}{2}})^{d+\alpha}} \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \overline{p}(t - \tilde{s}, x, w) m(dw) \int_{\mathbb{R}^d} \frac{1}{|x - z|^{d+\alpha}} \overline{p}(\tilde{s}, y, z) m(dy) d\tilde{s} \\
& \leq (\tilde{C}_1 (\overline{CM}^2 10^{d+\alpha}) C_t + \tilde{C}_1 (\overline{CM}^2 2^{\frac{1}{2}(d+\alpha)}) C_t \frac{t}{|x - z|^{d+\alpha}}) + \overline{C} \frac{L^2 D_2^2}{(1 - \frac{1}{10} - 2^{-\frac{1}{2}})^{d+\alpha}} \frac{t}{|x - z|^{d+\alpha}}.
\end{aligned}$$

There exists $t_{04} > 0$ such that when $0 < t \leq t_{04}$, C_t is small enough and the sum of the first two terms $\leq \frac{1}{4} \tilde{C}_1 K \frac{t}{|x - z|^{d+\alpha}}$. For the third term, we know that

$$\overline{C} \frac{L^2 D_2^2}{(1 - \frac{1}{10} - 2^{-\frac{1}{2}})^{d+\alpha}} \leq \frac{1}{4} \tilde{C}_1 K.$$

Thus

$$\int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \overline{p}(t - \tilde{s}, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{CF}^2(w, y)}{|w - y|^{d+\alpha}} \overline{p}(\tilde{s}, y, z) m(dy) d\tilde{s} \leq \frac{1}{2} \tilde{C}_1 K \frac{t}{|x - z|^{d+\alpha}}.$$

Combining case a and case b, when $0 < t \leq \min(t_{03}, t_{04})$,

$$\int_{\frac{t}{2}}^t \int_{\mathbb{R}^d} \overline{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{CF}^2(w, y)}{|w - y|^{d+\alpha}} \overline{p}(t - s, y, z) m(dy) ds \leq \frac{1}{2} \tilde{C}_1 K t^{-\frac{d}{\alpha}} \left(1 \wedge \frac{t^{\frac{1}{\alpha}}}{|x - z|} \right)^{d+\alpha}.$$

Therefore, when $0 < t \leq t_0 = \min(t_{01}, t_{02}, t_{03}, t_{04})$,

$$\overline{q}_1(t, x, z) \leq \tilde{C}_1 K t^{-\frac{d}{\alpha}} \left(1 \wedge \frac{t^{\frac{1}{\alpha}}}{|x - z|} \right)^{d+\alpha}.$$

i.e. statement 2 holds for $n = 1$.

Next we suppose statement 1 and statement 2 are true for $n \leq m - 1$, we show that they hold for $n = m$.

First we look at statement 1.

For the first part of statement 1,

$$\begin{aligned}
& \int_{\mathbb{R}^d} \overline{q}_m(t, x, z) m(dz) \\
& \leq \int_{\mathbb{R}^d} C_m^1 \int_0^t \int_{\mathbb{R}^d} \overline{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{CF}^2(w, y)}{|w - y|^{d+\alpha}} \overline{q}_{m-1}(t - s, y, z) m(dy) ds m(dz) \\
& \quad + \int_{\mathbb{R}^d} \sum_{i=2}^m C_m^i \int_0^t \int_{\mathbb{R}^d} \overline{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{CF}^2(w, y)}{|w - y|^{d+\alpha}} L^{i-2} \overline{q}_{m-i}(t - s, y, z) m(dy) ds m(dz) \\
& \leq C_m^1 \int_0^t \int_{\mathbb{R}^d} \overline{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{CF}^2(w, y)}{|w - y|^{d+\alpha}} \left(\int_{\mathbb{R}^d} \overline{q}_{m-1}(t - s, y, z) m(dz) \right) m(dy) ds
\end{aligned}$$

$$\begin{aligned}
& + \sum_{i=2}^{m-1} C_m^i \int_0^t \int_{\mathbb{R}^d} \bar{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\bar{C}^2 \bar{F}^2(w, y)}{|w - y|^{d+\alpha}} L^{i-2} \left(\int_{\mathbb{R}^d} \bar{q}_{m-i}(t - s, y, z) m(dz) \right) m(dy) ds \\
& + \int_0^t \int_{\mathbb{R}^d} \bar{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\bar{C}^2 \bar{F}^2(w, y)}{|w - y|^{d+\alpha}} L^{m-2} \left(\int_{\mathbb{R}^d} \bar{p}(t - s, y, z) m(dz) \right) m(dy) ds \\
& \leq C_m^1 \tilde{C} C_t (m-1)! K^{m-1} \int_0^t \int_{\mathbb{R}^d} \bar{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\bar{C} \bar{F}^2(w, y)}{|w - y|^{d+\alpha}} m(dy) ds \\
& + \sum_{i=2}^{m-1} C_m^i \tilde{C} C_t (m-i)! L^{i-2} K^{m-i} \int_0^t \int_{\mathbb{R}^d} \bar{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\bar{C}^2 \bar{F}^2(w, y)}{|w - y|^{d+\alpha}} m(dy) ds \\
& \quad (\text{since statement 1 holds for } n \leq m-1) \\
& + \int_0^t \int_{\mathbb{R}^d} \bar{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\bar{C}^2 \bar{F}^2(w, y)}{|w - y|^{d+\alpha}} L^{m-2} m(dy) ds D_2 \\
& \leq \tilde{C} C_t (m-1)! K^{m-1} \bar{C} \bar{M}^2 C_t + \sum_{i=2}^{m-1} C_m^i \tilde{C} C_t (m-i)! L^{i-2} K^{m-i} \bar{C}^2 \bar{M}^2 C_t + L^{m-2} \bar{C}^2 \bar{M}^2 D_2 C_t.
\end{aligned}$$

For the sum of the first and the second term, there exists $t_{11} > 0$ with $t_{11} \leq t_0$ such that when $0 < t < t_{11}$, C_t is small enough and

$$\tilde{C} C_t \left(C_m^1 (m-1)! K^{m-1} + \sum_{i=2}^{m-1} C_m^i (m-i)! L^{i-2} K^{m-i} \bar{C} \right) \bar{C} \bar{M}^2 \leq \frac{1}{2} \tilde{C} m! K^m.$$

For the third term, we know that

$$L^{m-2} \bar{C}^2 \bar{M}^2 D_2 \leq \frac{1}{2} \tilde{C} m! K^m, \quad \forall m > 0.$$

Thus

$$\int_{\mathbb{R}^d} \bar{q}_m(t, x, z) m(dz) \leq \tilde{C} C_t m! K^m.$$

i.e. the first part of the statement 1 holds for $n = m$.

To show the second part of the statement 1 for $n = m$, we know that

$$\begin{aligned}
& \bar{q}_m(t, x, z) \\
& \leq C_m^1 \int_0^t \int_{\mathbb{R}^d} \bar{p}(s_1, x, w_1) m(dw_1) \int_{\mathbb{R}^d} \frac{\bar{C} \bar{F}^2(w_1, y_1)}{|w_1 - y_1|^{d+\alpha}} \bar{q}_{m-1}(t - s_1, y_1, z) m(dy_1) ds_1 \\
& + \sum_{i=2}^m C_m^i \int_0^t \int_{\mathbb{R}^d} \bar{p}(s_1, x, w_1) m(dw_1) \int_{\mathbb{R}^d} \frac{\bar{C}^2 \bar{F}^2(w_1, y_1)}{|w_1 - y_1|^{d+\alpha}} L^{i-2} \bar{q}_{m-i}(t - s_1, y_1, z) m(dy_1) ds_1.
\end{aligned}$$

Substituting the expressions of $\bar{q}_{m-1}(t-s_1, y_1, z)$ and $\bar{q}_{m-i}(t-s_1, y_1, z)$ into the above, we have

$$\begin{aligned}
& \bar{q}_m(t, x, z) \\
& \leq C_m^1 \int_0^t \int_{\mathbb{R}^d} \bar{p}(s_1, x, w_1) m(dw_1) \int_{\mathbb{R}^d} \frac{\overline{CF}^2(w_1, y_1)}{|w_1 - y_1|^{d+\alpha}} m(dy_1) ds_1 (C_{m-1}^1 \int_0^{t-s_1} \int_{\mathbb{R}^d} \bar{p}(s_2, y_1, w_2) \\
& \quad \cdot m(dw_2) \int_{\mathbb{R}^d} \frac{\overline{CF}^2(w_2, y_2)}{|w_2 - y_2|^{d+\alpha}} \bar{q}_{m-2}(t-s_1-s_2, y_2, z) m(dy_2) ds_2) \\
& \quad + C_m^1 \int_0^t \int_{\mathbb{R}^d} \bar{p}(s_1, x, w_1) m(dw_1) \int_{\mathbb{R}^d} \frac{\overline{CF}^2(w_1, y_1)}{|w_1 - y_1|^{d+\alpha}} m(dy_1) ds_1 \left(\sum_{i=2}^{m-1} C_{m-1}^i \int_0^{t-s_1} \right. \\
& \quad \cdot \int_{\mathbb{R}^d} \bar{p}(s_2, y_1, w_2) m(dw_2) \int_{\mathbb{R}^d} \frac{\overline{C^2 F^2}(w_2, y_2)}{|w_2 - y_2|^{d+\alpha}} L^{i-2} \bar{q}_{m-1-i}(t-s_1-s_2, y_2, z) m(dy_2) ds_2) \\
& \quad + \sum_{i=2}^m C_m^i \int_0^t \int_{\mathbb{R}^d} \bar{p}(s_1, x, w_1) m(dw_1) \int_{\mathbb{R}^d} \frac{\overline{C^2 F^2}(w_1, y_1)}{|w_1 - y_1|^{d+\alpha}} L^{i-2} m(dy_1) ds_1 (C_{m-i}^1 \int_0^{t-s_1} \\
& \quad \cdot \int_{\mathbb{R}^d} \bar{p}(s_2, y_1, w_2) m(dw_2) \int_{\mathbb{R}^d} \frac{\overline{CF}^2(w_2, y_2)}{|w_2 - y_2|^{d+\alpha}} \bar{q}_{m-i-1}(t-s_1-s_2, y_2, z) m(dy_2) ds_2) \\
& \quad + \sum_{i=2}^m C_m^i \int_0^t \int_{\mathbb{R}^d} \bar{p}(s_1, x, w_1) m(dw_1) \int_{\mathbb{R}^d} \frac{\overline{C^2 F^2}(w_1, y_1)}{|w_1 - y_1|^{d+\alpha}} L^{i-2} m(dy_1) ds_1 \left(\sum_{j=2}^{m-i} C_{m-i}^j \int_0^{t-s_1} \right. \\
& \quad \cdot \int_{\mathbb{R}^d} \bar{p}(s_2, y_1, w_2) m(dw_2) \int_{\mathbb{R}^d} \frac{\overline{C^2 F^2}(w_2, y_2)}{|w_2 - y_2|^{d+\alpha}} L^{j-2} \bar{q}_{m-i-j}(t-s_1-s_2, y_2, z) m(dy_2) ds_2) \\
& \leq C_m^1 \int_0^t \int_{\mathbb{R}^d} \bar{p}(s_1, x, w_1) m(dw_1) \int_{\mathbb{R}^d} \frac{\overline{CF}^2(w_1, y_1)}{|w_1 - y_1|^{d+\alpha}} m(dy_1) ds_1 (C_{m-1}^1 \int_0^{t-s_1} \int_{\mathbb{R}^d} \bar{p}(s_2, y_1, w_2) \\
& \quad \cdot m(dw_2) \int_{\mathbb{R}^d} \frac{\overline{CF}^2(w_2, y_2)}{|w_2 - y_2|^{d+\alpha}} \tilde{C}_1(m-2)! K^{m-2} \frac{1}{2M_1} \bar{p}(t-s_1-s_2, y_2, z) m(dy_2) ds_2) \\
& \quad + C_m^1 \int_0^t \int_{\mathbb{R}^d} \bar{p}(s_1, x, w_1) m(dw_1) \int_{\mathbb{R}^d} \frac{\overline{CF}^2(w_1, y_1)}{|w_1 - y_1|^{d+\alpha}} m(dy_1) ds_1 \left(\sum_{i=2}^{m-1} C_{m-1}^i \int_0^{t-s_1} \int_{\mathbb{R}^d} \right. \\
& \quad \cdot \bar{p}(s_2, y_1, w_2) m(dw_2) \int_{\mathbb{R}^d} \frac{\overline{C^2 F^2}(w_2, y_2)}{|w_2 - y_2|^{d+\alpha}} \tilde{C}_1(m-1-i)! L^{i-2} K^{m-1-i} \frac{1}{2M_1} \\
& \quad \cdot \bar{p}(t-s_1-s_2, y_2, z) m(dy_2) ds_2) \\
& \quad + \sum_{i=2}^m C_m^i \int_0^t \int_{\mathbb{R}^d} \bar{p}(s_1, x, w_1) m(dw_1) \int_{\mathbb{R}^d} \frac{\overline{C^2 F^2}(w_1, y_1)}{|w_1 - y_1|^{d+\alpha}} L^{i-2} m(dy_1) ds_1 (C_{m-i}^1 \int_0^{t-s_1} \int_{\mathbb{R}^d} \\
& \quad \cdot \bar{p}(s_2, y_1, w_2) m(dw_2) \int_{\mathbb{R}^d} \frac{\overline{CF}^2(w_2, y_2)}{|w_2 - y_2|^{d+\alpha}} \tilde{C}_1(m-i-1)! K^{m-i-1} \frac{1}{2M_1} \\
& \quad \cdot \bar{p}(t-s_1-s_2, y_2, z) m(dy_2) ds_2) \\
& \quad + \sum_{i=2}^m C_m^i \int_0^t \int_{\mathbb{R}^d} \bar{p}(s_1, x, w_1) m(dw_1) \int_{\mathbb{R}^d} \frac{\overline{C^2 F^2}(w_1, y_1)}{|w_1 - y_1|^{d+\alpha}} L^{i-2} m(dy_1) ds_1 \left(\sum_{j=2}^{m-i} C_{m-i}^j \int_0^{t-s_1} \right.
\end{aligned}$$

$$\begin{aligned}
& \cdot \int_{\mathbb{R}^d} \bar{p}(s_2, y_1, w_2) m(dw_2) \int_{\mathbb{R}^d} \frac{\bar{C}^2 \bar{F}^2(w_2, y_2)}{|w_2 - y_2|^{d+\alpha}} \tilde{C}_1(m-i-j)! L^{j-2} K^{m-i-j} \frac{1}{2M_1} \\
& \cdot \bar{p}(t-s_1-s_2, y_2, z) m(dy_2) ds_2) \\
& (\text{ since the statement 2 holds true for } n \leq m-1) \\
& = C_m^1 C_{m-1}^1 \bar{C}^2 \tilde{C}_1(m-2)! K^{m-2} \frac{1}{2M_1} \left(\int_0^t \int_{\mathbb{R}^d} \bar{p}(s_1, x, w_1) m(dw_1) \int_{\mathbb{R}^d} \frac{\bar{F}^2(w_1, y_1)}{|w_1 - y_1|^{d+\alpha}} \right. \\
& \cdot m(dy_1) ds_1 \int_0^{t-s_1} \int_{\mathbb{R}^d} \bar{p}(s_2, y_1, w_2) m(dw_2) \int_{\mathbb{R}^d} \frac{\bar{F}^2(w_2, y_2)}{|w_2 - y_2|^{d+\alpha}} \bar{p}(t-s_1-s_2, y_2, z) m(dy_2) ds_2) \\
& + C_m^1 \sum_{i=2}^{m-1} C_{m-1}^i \bar{C}^3 \tilde{C}_1(m-1-i)! L^{i-2} K^{m-1-i} \frac{1}{2M_1} \left(\int_0^t \int_{\mathbb{R}^d} \bar{p}(s_1, x, w_1) m(dw_1) \right. \\
& \cdot \int_{\mathbb{R}^d} \frac{\bar{F}^2(w_1, y_1)}{|w_1 - y_1|^{d+\alpha}} m(dy_1) ds_1 \int_0^{t-s_1} \int_{\mathbb{R}^d} \bar{p}(s_2, y_1, w_2) m(dw_2) \int_{\mathbb{R}^d} \frac{\bar{F}^2(w_2, y_2)}{|w_2 - y_2|^{d+\alpha}} \\
& \cdot \bar{p}(t-s_1-s_2, y_2, z) m(dy_2) ds_2) \\
& + \sum_{i=2}^m C_m^i C_{m-i}^1 \bar{C}^3 \tilde{C}_1(m-i-1)! L^{i-2} K^{m-i-1} \frac{1}{2M_1} \left(\int_0^t \int_{\mathbb{R}^d} \bar{p}(s_1, x, w_1) m(dw_1) \right. \\
& \cdot \int_{\mathbb{R}^d} \frac{\bar{F}^2(w_1, y_1)}{|w_1 - y_1|^{d+\alpha}} m(dy_1) ds_1 \int_0^{t-s_1} \int_{\mathbb{R}^d} \bar{p}(s_2, y_1, w_2) m(dw_2) \int_{\mathbb{R}^d} \frac{\bar{F}^2(w_2, y_2)}{|w_2 - y_2|^{d+\alpha}} \\
& \cdot \bar{p}(t-s_1-s_2, y_2, z) m(dy_2) ds_2) \\
& + \sum_{i=2}^m C_m^i \sum_{j=2}^{m-i} C_{m-i}^j \bar{C}^4 \tilde{C}_1(m-i-j)! L^{i+j-4} K^{m-i-j} \frac{1}{2M_1} \left(\int_0^t \int_{\mathbb{R}^d} \bar{p}(s_1, x, w_1) m(dw_1) \right. \\
& \cdot \int_{\mathbb{R}^d} \frac{\bar{F}^2(w_1, y_1)}{|w_1 - y_1|^{d+\alpha}} m(dy_1) ds_1 \int_0^{t-s_1} \int_{\mathbb{R}^d} \bar{p}(s_2, y_1, w_2) m(dw_2) \int_{\mathbb{R}^d} \frac{\bar{F}^2(w_2, y_2)}{|w_2 - y_2|^{d+\alpha}} \\
& \cdot \bar{p}(t-s_1-s_2, y_2, z) m(dy_2) ds_2).
\end{aligned}$$

It is clear that in the above expression there is the following common factor :

$$\begin{aligned}
& \int_0^t \int_{\mathbb{R}^d} \bar{p}(s_1, x, w_1) m(dw_1) \int_{\mathbb{R}^d} \frac{\bar{F}^2(w_1, y_1)}{|w_1 - y_1|^{d+\alpha}} m(dy_1) ds_1 \int_0^{t-s_1} \int_{\mathbb{R}^d} \bar{p}(s_2, y_1, w_2) m(dw_2) \\
& \cdot \int_{\mathbb{R}^d} \frac{\bar{F}^2(w_2, y_2)}{|w_2 - y_2|^{d+\alpha}} \bar{p}(t-s_1-s_2, y_2, z) m(dy_2) ds_2.
\end{aligned}$$

We denote it by $G(t, x, z)$.

Next we show $G(t, x, z) = G(t, z, x)$.

Changing variables, let $t-s_1-s_2 = \tilde{s}_1$, $s_2 = \tilde{s}_2$. The absolute value of the Jacobian of the this transformation is 1. Let $y_2 = \tilde{w}_1$, $y_1 = \tilde{w}_2$, $w_2 = \tilde{y}_1$, $w_1 = \tilde{y}_2$.

Thus

$$\begin{aligned}
G(t, x, z) &= \int_0^t \int_0^{t-\tilde{s}_1} \int_{\mathbb{R}^d} \bar{p}(t - \tilde{s}_1 - \tilde{s}_2, x, \tilde{y}_2) m(d\tilde{y}_2) \int_{\mathbb{R}^d} \frac{\bar{F}^2(\tilde{y}_2, \tilde{w}_2)}{|\tilde{y}_2 - \tilde{w}_2|^{d+\alpha}} m(d\tilde{w}_2) \\
&\quad \cdot \left(\int_{\mathbb{R}^d} \bar{p}(\tilde{s}_2, \tilde{w}_2, \tilde{y}_1) m(d\tilde{y}_1) \int_{\mathbb{R}^d} \frac{\bar{F}^2(\tilde{y}_1, \tilde{w}_1)}{|\tilde{y}_1 - \tilde{w}_1|^{d+\alpha}} \bar{p}(\tilde{s}_1, \tilde{w}_1, z) m(d\tilde{w}_1) d\tilde{s}_2 d\tilde{s}_1. \right)
\end{aligned}$$

Rearranging the components of the integrand and using the fact that $\bar{F}(x, y)$ and $\bar{p}(t, x, y)$ are symmetric in x and y , we can see that the expression of $G(t, x, z)$ is equal to $G(t, z, x)$.

Therefore

$$\int_{\mathbb{R}^d} G(t, x, z) m(dx) = \int_{\mathbb{R}^d} G(t, z, x) m(dx),$$

and both

$$\begin{aligned}
&= \int_0^t \int_{\mathbb{R}^d} \bar{p}(s_1, z, w_1) m(dw_1) \int_{\mathbb{R}^d} \frac{\bar{F}^2(w_1, y_1)}{|w_1 - y_1|^{d+\alpha}} m(dy_1) ds_1 \left(\int_0^{t-s_1} \int_{\mathbb{R}^d} \bar{p}(s_2, y_1, w_2) m(dw_2) \right. \\
&\quad \cdot \left. \left(\int_0^{t-s_1} \int_{\mathbb{R}^d} \bar{p}(s_2, y_1, w_2) m(dw_2) \int_{\mathbb{R}^d} \frac{\bar{F}^2(w_2, y_2)}{|w_2 - y_2|^{d+\alpha}} \int_{\mathbb{R}^d} \bar{p}(t - s_1 - s_2, y_2, x) m(dx) m(dy_2) ds_2 \right) \right) \\
&= \int_0^t \int_{\mathbb{R}^d} \bar{p}(s_1, z, w_1) m(dw_1) \int_{\mathbb{R}^d} \frac{\bar{F}^2(w_1, y_1)}{|w_1 - y_1|^{d+\alpha}} D_2 m(dy_1) ds_1 \\
&\quad \cdot \left(\int_0^{t-s_1} \int_{\mathbb{R}^d} \bar{p}(s_2, y_1, w_2) m(dw_2) \int_{\mathbb{R}^d} \frac{\bar{F}^2(w_2, y_2)}{|w_2 - y_2|^{d+\alpha}} m(dy_2) ds_2 \right) \\
&\leq \int_0^t \int_{\mathbb{R}^d} \bar{p}(s_1, z, w_1) m(dw_1) \int_{\mathbb{R}^d} \frac{\bar{F}^2(w_1, y_1)}{|w_1 - y_1|^{d+\alpha}} m(dy_1) ds_1 \left(\int_0^{t-s_1} \int_{\mathbb{R}^d} \bar{p}(s_2, y_1, w_2) \mu(dw_2) ds_2 D_2 \bar{M}^2 \right) \\
&\leq \int_0^t \int_{\mathbb{R}^d} \bar{p}(s_1, z, w_1) m(dw_1) \int_{\mathbb{R}^d} \frac{\bar{F}^2(w_1, y_1)}{|w_1 - y_1|^{d+\alpha}} m(dy_1) ds_1 C_t D_2 \bar{M}^2 \\
&\leq \int_0^t \int_{\mathbb{R}^d} \bar{p}(s_1, z, w_1) \mu dw_1 \bar{M}^2 C_t D_2 \bar{M}^2 \leq D_2 \bar{M}^4 C_t C_t,
\end{aligned}$$

which implies that

$$\begin{aligned}
&\int_{\mathbb{R}^d} \bar{q}_m(t, x, z) m(dx) \\
&\leq C_m^1 C_{m-1}^1 \bar{C}^2 \tilde{C}_1 (m-2)! K^{m-2} \frac{1}{2M_1} D_2 \bar{M}^4 C_t C_t \\
&\quad + C_m^1 \sum_{i=2}^{m-1} C_{m-1}^i \bar{C}^3 \tilde{C}_1 (m-1-i)! L^{i-2} K^{m-1-i} \frac{1}{2M_1} D_2 \bar{M}^4 C_t C_t \\
&\quad + \sum_{i=2}^m C_m^i C_{m-i}^1 \bar{C}^3 \tilde{C}_1 (m-i-1)! L^{i-2} K^{m-i-1} \frac{1}{2M_1} D_2 \bar{M}^4 C_t C_t
\end{aligned}$$

$$+ \sum_{i=2}^m C_m^i \sum_{j=2}^{m-i} C_{m-i}^j \overline{C}^4 \tilde{C}_1 (m-i-j)! L^{i-2} L^{j-2} K^{m-i-j} \frac{1}{2M_1} D_2 \overline{M}^4 C_t C_t.$$

Since $C_t \downarrow 0$ as $t \downarrow 0$ and Lemma 3.1 and Lemma 3.2 hold, it is easy to see that there exists $t_{12} > 0$ with $t_{12} \leq t_0$ such that when $0 < t \leq t_{12}$, the first three term in the above expression $\leq \frac{1}{2} \tilde{C} C_t m! K^m$ and the fourth term $\leq \frac{1}{2} \tilde{C} C_t m! K^m$. Thus

$$\int_{\mathbb{R}^d} \overline{q}_m(t, x, z) m(dx) \leq \tilde{C} C_t m! K^m,$$

i.e. the second part of statement 1 holds for $n = m$. Thus statement 1 holds for $n = m$.

Next we show statement 2 for $n = m$. We write $\overline{q}_m(t, x, z)$ in the following way

$$\begin{aligned} & \overline{q}_m(t, x, z) \\ &= C_m^1 \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \overline{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{C} \overline{F}^2(w, y)}{|w - y|^{d+\alpha}} \overline{q}_{m-1}(t - s, y, z) m(dy) ds \\ &+ \sum_{i=2}^m C_m^i \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \overline{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{C}^i \overline{F}^i(w, y)}{|w - y|^{d+\alpha}} \overline{q}_{m-i}(t - s, y, z) m(dy) ds. \\ &+ C_m^1 \int_{\frac{t}{2}}^t \int_{\mathbb{R}^d} \overline{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{C} \overline{F}^2(w, y)}{|w - y|^{d+\alpha}} \overline{q}_{m-1}(t - s, y, z) m(dy) ds \\ &+ \sum_{i=2}^m C_m^i \int_{\frac{t}{2}}^t \int_{\mathbb{R}^d} \overline{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{C}^i \overline{F}^i(w, y)}{|w - y|^{d+\alpha}} \overline{q}_{m-i}(t - s, y, z) m(dy) ds. \end{aligned}$$

First we look at the sum of the first and the second terms in the expression of $\overline{q}_m(t, x, z)$.

There are two cases:

Case 1. When $|x - z| \leq t^{\frac{1}{\alpha}}$,

$$\begin{aligned} & C_m^1 \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \overline{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{C} \overline{F}^2(w, y)}{|w - y|^{d+\alpha}} \overline{q}_{m-1}(t - s, y, z) m(dy) ds \\ &+ \sum_{i=2}^m C_m^i \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \overline{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{C}^i \overline{F}^i(w, y)}{|w - y|^{d+\alpha}} \overline{q}_{m-i}(t - s, y, z) m(dy) ds. \\ &\leq C_m^1 \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \overline{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{F}^2(w, y)}{|w - y|^{d+\alpha}} dy ds \overline{C} \overline{M} \tilde{C}_1 (m-1)! K^{m-1} \left(\frac{t}{2}\right)^{-\frac{d}{\alpha}} \\ &+ \sum_{i=2}^m C_m^i \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \overline{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{F}^2(w, y)}{|w - y|^{d+\alpha}} dy ds \overline{C}^2 \overline{M} L^{i-2} \tilde{C}_1 (m-i)! K^{m-i} \left(\frac{t}{2}\right)^{-\frac{d}{\alpha}} \\ &\leq C_m^1 C_t \overline{C} \overline{M}^2 \tilde{C}_1 (m-1)! K^{m-1} \left(\frac{t}{2}\right)^{-\frac{d}{\alpha}} \end{aligned}$$

$$\begin{aligned}
& + \sum_{i=2}^m C_m^i C_t \overline{C}^2 \overline{M}^2 \tilde{C}_1 (m-i)! L^{i-2} K^{m-i} \left(\frac{t}{2}\right)^{-\frac{d}{\alpha}} \\
& = \tilde{C}_1 C_t \overline{C} \overline{M}^2 \left(C_m^1 (m-1)! K^{m-1} + \overline{C} \sum_{i=2}^m C_m^i (m-i)! L^{i-2} K^{m-i} \right) \left(\frac{t}{2}\right)^{-\frac{d}{\alpha}}.
\end{aligned}$$

There exists $t_{13} > 0$ with $t_{13} \leq t_0$ such that when $0 < t \leq t_{13}$, C_t is small enough,

$$C_t \overline{C} \overline{M}^2 \left(C_m^1 (m-1)! K^{m-1} + \overline{C} \sum_{i=2}^m C_m^i (m-i)! L^{i-2} K^{m-i} \right) \left(\frac{1}{2}\right)^{-\frac{d}{\alpha}} \leq \frac{1}{2} m! K^m.$$

Thus

$$\begin{aligned}
& C_m^1 \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \overline{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{C} \overline{F}^2(w, y)}{|w-y|^{d+\alpha}} \overline{q}_{m-1}(t-s, y, z) m(dy) ds \\
& + \sum_{i=2}^m C_m^i \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \overline{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{C}^i \overline{F}^i(w, y)}{|w-y|^{d+\alpha}} \overline{q}_{m-i}(t-s, y, z) m(dy) ds \\
& \leq \frac{1}{2} \tilde{C}_1 m! K^m t^{-\frac{d}{\alpha}}.
\end{aligned}$$

Case 2. When $|x-z| \geq t^{\frac{1}{\alpha}}$. Let $B_1 = \{y \in \mathbb{R}^d \mid |y-z| \geq \frac{1}{10}|x-z|\}$, $B_2 = \{w \in \mathbb{R}^d \mid |w-x| \geq 2^{-\frac{1}{2}}|x-z|\}$ and $B_3 = \{(w, y) \in \mathbb{R}^d \times \mathbb{R}^d \mid |y-z| < \frac{1}{10}|x-z|, |w-x| < 2^{-\frac{1}{2}}|x-z|\}$. On B_3 , we have $|w-y| \geq (1 - \frac{1}{10} - 2^{-\frac{1}{2}})|x-z|$.

$$\begin{aligned}
& C_m^1 \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \overline{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{C} \overline{F}^2(w, y)}{|w-y|^{d+\alpha}} \overline{q}_{m-1}(t-s, y, z) m(dy) ds \\
& + \sum_{i=2}^m C_m^i \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \overline{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{C}^i \overline{F}^i(w, y)}{|w-y|^{d+\alpha}} \overline{q}_{m-i}(t-s, y, z) m(dy) ds \\
& \leq C_m^1 \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \overline{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{C} \overline{F}^2(w, y)}{|w-y|^{d+\alpha}} \overline{q}_{m-1}(t-s, y, z) 1_{B_1}(y) m(dy) ds \\
& + \sum_{i=2}^m C_m^i \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \overline{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{C}^i \overline{F}^i(w, y)}{|w-y|^{d+\alpha}} \overline{q}_{m-i}(t-s, y, z) 1_{B_1}(y) m(dy) ds \\
& + C_m^1 \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \overline{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{C} \overline{F}^2(w, y)}{|w-y|^{d+\alpha}} \overline{q}_{m-1}(t-s, y, z) 1_{B_2}(w) m(dy) ds \\
& + \sum_{i=2}^m C_m^i \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \overline{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{C}^i \overline{F}^i(w, y)}{|w-y|^{d+\alpha}} \overline{q}_{m-i}(t-s, y, z) 1_{B_2}(w) m(dy) ds
\end{aligned}$$

$$\begin{aligned}
& + C_m^1 \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \bar{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{CF}^2(w, y)}{|w - y|^{d+\alpha}} \bar{q}_{m-1}(t - s, y, z) 1_{B_3}(w, y) m(dy) ds \\
& + \sum_{i=2}^m C_m^i \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \bar{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{C^i F^i}(w, y)}{|w - y|^{d+\alpha}} \bar{q}_{m-i}(t - s, y, z) 1_{B_3}(w, y) m(dy) ds \\
& \leq C_m^1 \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \bar{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{F}^2(w, y)}{|w - y|^{d+\alpha}} \overline{MC} \tilde{C}_1(m-1)! K^{m-1} 10^{d+\alpha} \frac{(t-s)}{|x-z|^{d+\alpha}} dy ds \\
& + \sum_{i=2}^m C_m^i \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \bar{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{F}^2(w, y)}{|w - y|^{d+\alpha}} \overline{MC}^2 L^{i-2} \tilde{C}_1(m-i)! K^{m-i} \frac{(t-s)}{|x-z|^{d+\alpha}} dy ds \\
& + C_m^1 \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} dw \tilde{C}_1 \overline{M}^2 2^{\frac{1}{2}(d+\alpha)} \frac{s}{|x-z|^{d+\alpha}} \int_{\mathbb{R}^d} \frac{\overline{F}^2(w, y)}{|w - y|^{d+\alpha}} \bar{q}_{m-1}(t-s, y, z) dy ds \overline{C} \\
& + \sum_{i=2}^m C_m^i \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} dw \tilde{C}_1 \overline{M}^2 2^{\frac{1}{2}(d+\alpha)} \frac{s}{|x-z|^{d+\alpha}} \int_{\mathbb{R}^d} \frac{\overline{F}^2(w, y)}{|w - y|^{d+\alpha}} \bar{q}_{m-i}(t-s, y, z) dy ds \overline{C}^2 L^{i-2} \\
& + C_m^1 \overline{C} \frac{L^2}{(1 - \frac{1}{10} - 2^{-\frac{1}{2}})^{d+\alpha}} \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \bar{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{1}{|x-z|^{d+\alpha}} \bar{q}_{m-1}(t-s, y, z) m(dy) ds \\
& + \sum_{i=2}^{m-1} C_m^i \frac{L^i}{(1 - \frac{1}{10} - 2^{-\frac{1}{2}})^{d+\alpha}} \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \bar{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{1}{|x-z|^{d+\alpha}} \bar{q}_{m-i}(t-s, y, z) m(dy) ds \\
& + \frac{L^m}{(1 - \frac{1}{10} - 2^{-\frac{1}{2}})^{d+\alpha}} \frac{1}{|x-z|^{d+\alpha}} \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \bar{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \bar{p}(t-s, y, z) m(dy) ds \\
& \leq C_m^1 C_t \overline{CM}^2 \tilde{C}_1(m-1)! K^{m-1} 10^{d+\alpha} \frac{t}{|x-z|^{d+\alpha}} \\
& + \sum_{i=2}^m C_m^i C_t \overline{C}^2 \overline{M}^2 L^{i-2} \tilde{C}_1(m-i)! K^{m-i} 10^{d+\alpha} \frac{t}{|x-z|^{d+\alpha}} \\
& + C_m^1 \tilde{C}_1 \overline{M}^2 2^{\frac{1}{2}(d+\alpha)} \frac{\tilde{C}_1(m-1)! K^{m-1} t}{|x-z|^{d+\alpha}} \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} dw \int_{\mathbb{R}^d} \frac{\overline{F}^2(w, y)}{|w - y|^{d+\alpha}} \frac{1}{2M_1} \bar{p}(t-s, z, y) dy ds \overline{C} \\
& \quad (\text{by (3.3) in statement 1 for } n \leq m-1) \\
& + \sum_{i=2}^m C_m^i \tilde{C}_1 \overline{M}^2 2^{\frac{1}{2}(d+\alpha)} \frac{\tilde{C}_1(m-i)! K^{m-i} t}{|x-z|^{d+\alpha}} \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} dw \int_{\mathbb{R}^d} \frac{\overline{F}^2(w, y)}{|w - y|^{d+\alpha}} \frac{1}{2M_1} \bar{p}(t-s, z, y) dy ds \\
& \cdot \overline{C}^2 L^{i-2} + C_m^1 \frac{\overline{C} L^2}{(1 - \frac{1}{10} - 2^{-\frac{1}{2}})^{d+\alpha}} \frac{\frac{t}{2}}{|x-z|^{d+\alpha}} D_2 \tilde{C} C_t(m-1)! K^{m-1} \\
& \quad (\text{by (3.4) in statement 1 for } n \leq m-1 \text{ and } \int_{\mathbb{R}^d} \bar{p}(s, x, w) m(dw) \leq D_2) \\
& + \sum_{i=2}^{m-1} C_m^i \frac{L^i}{(1 - \frac{1}{10} - 2^{-\frac{1}{2}})^{d+\alpha}} \frac{\frac{t}{2}}{|x-z|^{d+\alpha}} D_2 \tilde{C} C_t(m-i)! K^{m-i} \\
& + \frac{L^m}{(1 - \frac{1}{10} - 2^{-\frac{1}{2}})^{d+\alpha}} \frac{t}{|x-z|^{d+\alpha}} D_2^2
\end{aligned}$$

$$\begin{aligned}
&\leq C_m^1 C_t \overline{C} \overline{M}^2 \tilde{C}_1 (m-1)! K^{m-1} 10^{d+\alpha} \frac{t}{|x-z|^{d+\alpha}} \\
&\quad + \sum_{i=2}^m C_m^i C_t \overline{C}^2 \overline{M}^2 (m-i)! \tilde{C}_1 L^{i-2} K^{m-i} 10^{d+\alpha} \frac{t}{|x-z|^{d+\alpha}} \\
&\quad + C_m^1 \tilde{C}_1 \overline{C} \overline{M}^2 2^{\frac{1}{2}(d+\alpha)} \frac{t}{|x-z|^{d+\alpha}} \tilde{C}_1 (m-1)! K^{m-1} \frac{1}{2M_1} C_t \\
&\quad + \sum_{i=2}^m C_m^i \tilde{C}_1 \overline{C}^2 \overline{M}^2 2^{\frac{1}{2}(d+\alpha)} \frac{t}{|x-z|^{d+\alpha}} \tilde{C}_1 (m-i)! K^{m-i} \frac{1}{2M_1} C_t L^{i-2} \\
&\quad + C_m^1 \overline{C} \frac{L^2}{(1 - \frac{1}{10} - 2^{-\frac{1}{2}})^{d+\alpha}} \frac{t}{|x-z|^{d+\alpha}} D_2 \tilde{C} C_t (m-1)! K^{m-1} \\
&\quad + \sum_{i=2}^{m-1} C_m^i \frac{L^i}{(1 - \frac{1}{10} - 2^{-\frac{1}{2}})^{d+\alpha}} \frac{t}{|x-z|^{d+\alpha}} D_2 \tilde{C} C_t (m-i)! K^{m-i} \\
&\quad + \frac{L^m D_2^2}{(1 - \frac{1}{10} - 2^{-\frac{1}{2}})^{d+\alpha}} \frac{t}{|x-z|^{d+\alpha}} \\
&\leq (C_m^1 \overline{C} \overline{M}^2 \tilde{C}_1 (m-1)! K^{m-1} 10^{d+\alpha}) C_t \frac{t}{|x-z|^{d+\alpha}} \\
&\quad + (\sum_{i=2}^m C_m^i \overline{C}^2 \overline{M}^2 (m-i)! \tilde{C}_1 L^{i-2} K^{m-i} 10^{d+\alpha}) C_t \frac{t}{|x-z|^{d+\alpha}} \\
&\quad + (C_m^1 \tilde{C}_1 \overline{C} \overline{M}^2 2^{\frac{1}{2}(d+\alpha)} \tilde{C}_1 (m-1)! K^{m-1} \frac{1}{2M_1}) C_t \frac{t}{|x-z|^{d+\alpha}} \\
&\quad + (\sum_{i=2}^m C_m^i \tilde{C}_1 \overline{C}^2 \overline{M}^2 2^{\frac{1}{2}(d+\alpha)} \tilde{C}_1 (m-i)! K^{m-i} \frac{1}{2M_1}) \frac{t}{|x-z|^{d+\alpha}} C_t L^{i-2} \\
&\quad + (C_m^1 \overline{C} \frac{L^2 D_2}{(1 - \frac{1}{10} - 2^{-\frac{1}{2}})^{d+\alpha}} \tilde{C} (m-1)! K^{m-1}) C_t \frac{t}{|x-z|^{d+\alpha}} \\
&\quad + (\sum_{i=2}^{m-1} C_m^i \overline{C} \frac{L^i D_2}{(1 - \frac{1}{10} - 2^{-\frac{1}{2}})^{d+\alpha}} \tilde{C} (m-i)! K^{m-i}) C_t \frac{t}{|x-z|^{d+\alpha}} \\
&\quad + \overline{C} \frac{L^m D_2^2}{(1 - \frac{1}{10} - 2^{-\frac{1}{2}})^{d+\alpha}} \frac{t}{|x-z|^{d+\alpha}}
\end{aligned}$$

There exists $t_{14} > 0$ with $t_{14} \leq t_0$ such that when $0 < t < t_{14}$, C_t is small enough and the sum of the first six terms in the above $\leq \frac{1}{4} \tilde{C}_1 m! K^m \frac{t}{|x-z|^{d+\alpha}}$. For the seventh term, we know that

$$\overline{C} \frac{L^m}{(1 - \frac{1}{10} - 2^{-\frac{1}{2}})^{d+\alpha}} \leq \frac{1}{4} \tilde{C}_1 m! K^m, \quad \forall m > 0.$$

Thus

$$\begin{aligned}
& C_m^1 \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \bar{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{CF}^2(w, y)}{|w - y|^{d+\alpha}} \bar{q}_{m-1}(t - s, y, z) m(dy) ds \\
& + \sum_{i=2}^m C_m^i \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \bar{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{CF}^i(w, y)}{|w - y|^{d+\alpha}} \bar{q}_{m-i}(t - s, y, z) m(dy) ds \\
& \leq \frac{1}{2} \tilde{C}_1 m! K^m \frac{t}{|x - z|^{d+\alpha}}.
\end{aligned}$$

Combining case 1 and case 2, when $0 < t \leq t_1 = \min(t_{11}, t_{12}, t_{13}, t_{14})$, the sum of the first and the second terms in the expression of $\bar{q}_m(t, x, z)$

$$\begin{aligned}
& C_m^1 \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \bar{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{CF}^2(w, y)}{|w - y|^{d+\alpha}} \bar{q}_{m-1}(t - s, y, z) m(dy) ds \\
& + \sum_{i=2}^m C_m^i \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \bar{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{CF}^i(w, y)}{|w - y|^{d+\alpha}} \bar{q}_{m-i}(t - s, y, z) m(dy) ds \\
& \leq \frac{1}{2} \tilde{C}_1 m! K^m t^{-\frac{d}{\alpha}} \left(1 \wedge \frac{t^{\frac{1}{\alpha}}}{|x - z|} \right)^{d+\alpha}.
\end{aligned}$$

For the sum of the third and the fourth terms in the expression of $\bar{q}_m(t, x, z)$

$$\begin{aligned}
& C_m^1 \int_{\frac{t}{2}}^t \int_{\mathbb{R}^d} \bar{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{CF}^2(w, y)}{|w - y|^{d+\alpha}} \bar{q}_{m-1}(t - s, y, z) m(dy) ds \\
& + \sum_{i=2}^m C_m^i \int_{\frac{t}{2}}^t \int_{\mathbb{R}^d} \bar{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{CF}^i(w, y)}{|w - y|^{d+\alpha}} \bar{q}_{m-i}(t - s, y, z) m(dy) ds,
\end{aligned}$$

let $t - s = \tilde{s}$. This sum becomes

$$\begin{aligned}
& C_m^1 \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \bar{p}(t - \tilde{s}, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{CF}^2(w, y)}{|w - y|^{d+\alpha}} \bar{q}_{m-1}(\tilde{s}, y, z) m(dy) d\tilde{s} \\
& + \sum_{i=2}^m C_m^i \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \bar{p}(t - \tilde{s}, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{CF}^i(w, y)}{|w - y|^{d+\alpha}} \bar{q}_{m-i}(\tilde{s}, y, z) m(dy) d\tilde{s}.
\end{aligned}$$

There are two cases:

Case a. When $|x - z| \leq t^{\frac{1}{\alpha}}$,

$$\begin{aligned}
& C_m^1 \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \bar{p}(t - \tilde{s}, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{CF}^2(w, y)}{|w - y|^{d+\alpha}} \bar{q}_{m-1}(\tilde{s}, y, z) m(dy) d\tilde{s} \\
& + \sum_{i=2}^m C_m^i \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \bar{p}(t - \tilde{s}, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{C^i F^i}(w, y)}{|w - y|^{d+\alpha}} \bar{q}_{m-i}(\tilde{s}, y, z) m(dy) d\tilde{s} \\
& \leq C_m^1 \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \tilde{C}_1 \left(\frac{t}{2}\right)^{-\frac{d}{\alpha}} m(dw) \int_{\mathbb{R}^d} \frac{\overline{F}^2(w, y)}{|w - y|^{d+\alpha}} \overline{C} \tilde{C}_1 (m-1)! K^{m-1} \frac{\bar{p}(\tilde{s}, z, y)}{M_1} m(dy) d\tilde{s} \\
& + \sum_{i=2}^m C_m^i \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \tilde{C}_1 \left(\frac{t}{2}\right)^{-\frac{d}{\alpha}} m(dw) \int_{\mathbb{R}^d} \frac{\overline{C^2 F^i}(w, y)}{|w - y|^{d+\alpha}} \tilde{C}_1 (m-i)! L^{i-2} K^{m-i} \frac{\bar{p}(\tilde{s}, z, y)}{M_1} m(dy) d\tilde{s} \\
& \leq C_m^1 \tilde{C}_1 \left(\frac{t}{2}\right)^{-\frac{d}{\alpha}} \overline{CM}^2 \tilde{C}_1 (m-1)! K^{m-1} \frac{1}{M_1} C_t + \sum_{i=2}^m C_m^i \tilde{C}_1 \left(\frac{t}{2}\right)^{-\frac{d}{\alpha}} \overline{C^2 M}^2 \tilde{C}_1 (m-i)! K^{m-i} \frac{1}{M_1} C_t \\
& + \sum_{i=2}^m C_m^i \tilde{C}_1 \left(\frac{t}{2}\right)^{-\frac{d}{\alpha}} \overline{C^2 M}^2 \tilde{C}_1 (m-i)! K^{m-i} \frac{1}{M_1} C_t \\
& = \tilde{C}_1 \overline{CM}^2 \left(\frac{t}{2}\right)^{-\frac{d}{\alpha}} \left(C_m^1 (m-1)! K^{m-1} \frac{1}{M_1} + \overline{C} \sum_{i=2}^m C_m^i (m-i)! K^{m-i} \frac{1}{M_1} \right) C_t
\end{aligned}$$

There exists $t_{21} > 0$ with $t_{21} \leq t_1$ such that when $0 < t \leq t_{21}$, C_t is small enough

$$\tilde{C}_1 \overline{CM}^2 \left(\frac{1}{2}\right)^{-\frac{d}{\alpha}} \left(C_m^1 (m-1)! K^{m-1} \frac{1}{M_1} + \overline{C} \sum_{i=2}^m C_m^i (m-i)! K^{m-i} \frac{1}{M_1} \right) \left(\frac{1}{2}\right)^{-\frac{d}{\alpha}} C_t \leq \frac{1}{2} \tilde{C}_1 m! K^m.$$

Thus

$$\begin{aligned}
& C_m^1 \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \bar{p}(t - \tilde{s}, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{CF}^2(w, y)}{|w - y|^{d+\alpha}} \bar{q}_{m-1}(\tilde{s}, y, z) m(dy) d\tilde{s} \\
& + \sum_{i=2}^m C_m^i \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \bar{p}(t - \tilde{s}, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{C^i F^i}(w, y)}{|w - y|^{d+\alpha}} \bar{q}_{m-i}(\tilde{s}, y, z) m(dy) d\tilde{s} \\
& \leq \frac{1}{2} \tilde{C}_1 m! K^m t^{-\frac{d}{\alpha}}.
\end{aligned}$$

Case b. When $|x - z| \geq t^{\frac{1}{\alpha}}$. Let $B_1 = \{y \in \mathbb{R}^d \mid |y - z| \geq \frac{1}{10}|x - z|\}$, $B_2 = \{w \in \mathbb{R}^d \mid |w - x| \geq 2^{-\frac{1}{2}}|x - z|\}$ and $B_3 = \{(w, y) \in \mathbb{R}^d \times \mathbb{R}^d \mid |y - z| < \frac{1}{10}|x - z|, |w - x| < 2^{-\frac{1}{2}}|x - z|\}$. On B_3 , we have $|w - y| \geq (1 - \frac{1}{10} - 2^{-\frac{1}{2}})|x - z|$.

$$\begin{aligned}
& C_m^1 \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \bar{p}(t - \tilde{s}, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{CF}^2(w, y)}{|w - y|^{d+\alpha}} \bar{q}_{m-1}(\tilde{s}, y, z) m(dy) d\tilde{s} \\
& + \sum_{i=2}^m C_m^i \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \bar{p}(t - \tilde{s}, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{C^i F^i}(w, y)}{|w - y|^{d+\alpha}} \bar{q}_{m-i}(\tilde{s}, y, z) m(dy) d\tilde{s} \\
& \leq C_m^1 \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \bar{p}(t - \tilde{s}, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{CF}^2(w, y)}{|w - y|^{d+\alpha}} \bar{q}_{m-1}(\tilde{s}, y, z) 1_{B_1}(y) m(dy) d\tilde{s} \\
& + \sum_{i=2}^m C_m^i \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \bar{p}(t - \tilde{s}, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{C^i F^i}(w, y)}{|w - y|^{d+\alpha}} \bar{q}_{m-i}(\tilde{s}, y, z) 1_{B_1}(y) m(dy) d\tilde{s} \\
& + C_m^1 \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \bar{p}(t - \tilde{s}, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{CF}^2(w, y)}{|w - y|^{d+\alpha}} \bar{q}_{m-1}(\tilde{s}, y, z) 1_{B_2}(w) m(dy) d\tilde{s} \\
& + \sum_{i=2}^m C_m^i \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \bar{p}(t - \tilde{s}, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{C^i F^i}(w, y)}{|w - y|^{d+\alpha}} \bar{q}_{m-i}(\tilde{s}, y, z) 1_{B_2}(w) m(dy) d\tilde{s} \\
& + C_m^1 \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \bar{p}(t - \tilde{s}, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{CF}^2(w, y)}{|w - y|^{d+\alpha}} \bar{q}_{m-1}(\tilde{s}, y, z) 1_{B_3}(w, y) m(dy) d\tilde{s} \\
& + \sum_{i=2}^m C_m^i \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \bar{p}(t - \tilde{s}, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{C^i F^i}(w, y)}{|w - y|^{d+\alpha}} \bar{q}_{m-i}(\tilde{s}, y, z) 1_{B_3}(w, y) m(dy) d\tilde{s} \\
& \leq C_m^1 \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \bar{p}(t - \tilde{s}, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{F}^2(w, y)}{|w - y|^{d+\alpha}} \overline{C} \tilde{C}_1(m-1)! K^{m-1} 10^{d+\alpha} \frac{\tilde{s}}{|x - z|^{d+\alpha}} m(dy) d\tilde{s} \\
& + \sum_{i=2}^m C_m^i \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \bar{p}(t - \tilde{s}, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{C^2 F^2}(w, y)}{|w - y|^{d+\alpha}} \tilde{C}_1(m-i)! L^{i-2} K^{m-i} 10^{d+\alpha} \frac{\tilde{s}}{|x - z|^{d+\alpha}} \\
& \cdot m(dy) d\tilde{s} \\
& + C_m^1 \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} dw \tilde{C}_1 \overline{M}^2 2^{\frac{1}{2}(d+\alpha)} \frac{t - \tilde{s}}{|x - z|^{d+\alpha}} \int_{\mathbb{R}^d} \frac{\overline{F}^2(w, y)}{|w - y|^{d+\alpha}} \overline{C} \tilde{C}_1(m-1)! K^{m-1} \frac{\bar{p}(\tilde{s}, z, y)}{M_1} dy d\tilde{s} \\
& + \sum_{i=2}^m C_m^i \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} dw \tilde{C}_1 \overline{M}^2 2^{\frac{1}{2}(d+\alpha)} \frac{t - \tilde{s}}{|x - z|^{d+\alpha}} \int_{\mathbb{R}^d} \frac{\overline{C^2 F^2}(w, y)}{|w - y|^{d+\alpha}} \tilde{C}_1 L^{i-2} (m-i)! K^{m-i} \\
& \cdot \frac{\bar{p}(\tilde{s}, z, y)}{M_1} dy d\tilde{s} \\
& + C_m^1 \overline{C} \frac{L^2}{(1 - \frac{1}{10} - 2^{-\frac{1}{2}})^{d+\alpha}} \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \bar{p}(t - \tilde{s}, x, w) m(dw) \int_{\mathbb{R}^d} \frac{1}{|x - z|^{d+\alpha}} \bar{q}_{m-1}(\tilde{s}, y, z) m(dy) d\tilde{s} \\
& + \sum_{i=2}^{m-1} C_m^i \frac{L^i}{(1 - \frac{1}{10} - 2^{-\frac{1}{2}})^{d+\alpha}} \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \bar{p}(t - \tilde{s}, x, w) m(dw) \int_{\mathbb{R}^d} \frac{1}{|x - z|^{d+\alpha}} \bar{q}_{m-i}(\tilde{s}, y, z) m(dy) \\
& \cdot d\tilde{s} + \frac{L^m}{(1 - \frac{1}{10} - 2^{-\frac{1}{2}})^{d+\alpha}} \frac{1}{|x - z|^{d+\alpha}} \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \bar{p}(t - \tilde{s}, x, w) m(dw) \int_{\mathbb{R}^d} \bar{p}(\tilde{s}, y, z) m(dy) d\tilde{s}
\end{aligned}$$

$$\begin{aligned}
&\leq C_m^1 C_t \overline{CM}^2 \tilde{C}_1 (m-1)! K^{m-1} 10^{d+\alpha} \frac{t}{|x-z|^{d+\alpha}} \\
&\quad + \sum_{i=2}^m C_m^i C_t \overline{C}^2 \overline{M}^2 \tilde{C}_1 (m-i)! K^{m-i} 10^{d+\alpha} \frac{t}{|x-z|^{d+\alpha}} \\
&\quad + C_m^1 \tilde{C}_1 \overline{CM}^2 2^{\frac{1}{2}(d+\alpha)} \frac{t}{|x-z|^{d+\alpha}} \tilde{C}_1 (m-1)! K^{m-1} \frac{1}{2M_1} C_t \\
&\quad + \sum_{i=2}^m C_m^i \tilde{C}_1 \overline{C}^2 \overline{M}^2 2^{\frac{1}{2}(d+\alpha)} \frac{t}{|x-z|^{d+\alpha}} \tilde{C}_1 (m-i)! L^{i-2} K^{m-i} \frac{1}{2M_1} C_t \\
&\quad + C_m^1 \frac{\overline{C} L^2 D_2}{(1 - \frac{1}{10} - 2^{-\frac{1}{2}})^{d+\alpha}} \frac{t}{|x-z|^{d+\alpha}} \tilde{C} C_t (m-1)! K^{m-1} \\
&\quad + \sum_{i=2}^{m-1} C_m^i \frac{L^i D_2}{(1 - \frac{1}{10} - 2^{-\frac{1}{2}})^{d+\alpha}} \frac{t}{|x-z|^{d+\alpha}} \tilde{C} C_t (m-i)! K^{m-i} \\
&\quad (\text{since statement 1 holds true for } n \leq m-1) \\
&\quad + \frac{L^m D_2^2}{(1 - \frac{1}{10} - 2^{-\frac{1}{2}})^{d+\alpha}} \frac{t}{|x-z|^{d+\alpha}} \\
&= (C_m^1 \overline{CM}^2 \tilde{C}_1 (m-1)! K^{m-1} 10^{d+\alpha}) C_t \frac{t}{|x-z|^{d+\alpha}} \\
&\quad + (\sum_{i=2}^m C_m^i \overline{C}^2 \overline{M}^2 \tilde{C}_1 (m-i)! K^{m-i} 10^{d+\alpha}) C_t \frac{t}{|x-z|^{d+\alpha}} \\
&\quad + (C_m^1 \tilde{C}_1 \overline{CM}^2 2^{\frac{1}{2}(d+\alpha)} \tilde{C}_1 (m-1)! K^{m-1} \frac{1}{2M_1}) C_t \frac{t}{|x-z|^{d+\alpha}} \\
&\quad + (\sum_{i=2}^m C_m^i \tilde{C}_1 \overline{C}^2 \overline{M}^2 2^{\frac{1}{2}(d+\alpha)} \tilde{C}_1 (m-i)! L^{i-2} K^{m-i} \frac{1}{2M_1}) C_t \frac{t}{|x-z|^{d+\alpha}} \\
&\quad + (C_m^1 \overline{C} \frac{L^2 D_2}{(1 - \frac{1}{10} - 2^{-\frac{1}{2}})^{d+\alpha}} \tilde{C} (m-1)! K^{m-1}) C_t \frac{t}{|x-z|^{d+\alpha}} \\
&\quad + (\sum_{i=2}^{m-1} C_m^i \frac{L^i D_2}{(1 - \frac{1}{10} - 2^{-\frac{1}{2}})^{d+\alpha}} \tilde{C} (m-i)! K^{m-i}) C_t \frac{t}{|x-z|^{d+\alpha}} \\
&\quad + \overline{C} \frac{L^m D_2^2}{(1 - \frac{1}{10} - 2^{-\frac{1}{2}})^{d+\alpha}} \frac{t}{|x-z|^{d+\alpha}}.
\end{aligned}$$

There exists $t_{22} > 0$ with $t_{21} \leq t_1$ such that when $0 < t \leq t_{22}$, C_t is small enough and the sum of the first six terms $\leq \frac{1}{4} \tilde{C}_1 m! K^m \frac{t}{|x-z|^{d+\alpha}}$. For the seventh term, we know that

$$\overline{C} \frac{L^m D_2^2}{(1 - \frac{1}{10} - 2^{-\frac{1}{2}})^{d+\alpha}} \leq \frac{1}{4} \tilde{C}_1 m! K^m.$$

Thus

$$\begin{aligned}
& C_m^1 \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \bar{p}(t - \tilde{s}, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{CF}^2(w, y)}{|w - y|^{d+\alpha}} \bar{q}_{m-1}(\tilde{s}, y, z) m(dy) d\tilde{s} \\
& + \sum_{i=2}^m C_m^i \int_0^{\frac{t}{2}} \int_{\mathbb{R}^d} \bar{p}(t - \tilde{s}, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{C^i F^i}(w, y)}{|w - y|^{d+\alpha}} \bar{q}_{m-i}(\tilde{s}, y, z) m(dy) d\tilde{s} \\
& \leq \frac{1}{2} \tilde{C}_1 m! K^m \frac{t}{|x - z|^{d+\alpha}}.
\end{aligned}$$

Combining case a and case b, when $0 < t < t_2 = \min(t_{21}, t_{22})$, the sum of the third and fourth terms of $\bar{q}(t, x, z)$

$$\begin{aligned}
& C_m^1 \int_{\frac{t}{2}}^t \int_{\mathbb{R}^d} \bar{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{CF}^2(w, y)}{|w - y|^{d+\alpha}} \bar{q}_{m-1}(t - s, y, z) m(dy) ds \\
& + \sum_{i=2}^m C_m^i \int_{\frac{t}{2}}^t \int_{\mathbb{R}^d} \bar{p}(s, x, w) m(dw) \int_{\mathbb{R}^d} \frac{\overline{C^i F^i}(w, y)}{|w - y|^{d+\alpha}} \bar{q}_{m-i}(t - s, y, z) m(dy) ds \\
& \leq \frac{1}{2} \tilde{C}_1 m! K^m t^{-\frac{d}{\alpha}} \left(1 \wedge \frac{t^{\frac{1}{\alpha}}}{|x - z|} \right)^{d+\alpha}.
\end{aligned}$$

Therefore, when $0 < t \leq t_2$,

$$\bar{q}_m(t, x, z) \leq \tilde{C}_1 m! K^m t^{-\frac{d}{\alpha}} \left(1 \wedge \frac{t^{\frac{1}{\alpha}}}{|x - z|} \right)^{d+\alpha},$$

i.e. the statement 2 holds for $n = m$. □

We know that for given $l > 0$, $q_n(t, x, z) = q_n^{(l)}(t, x, z)$, because of omission of the index l . However the expression of $\bar{q}_n(t, x, z)$ doesn't depend on l . So the constant t_2 in Theorem doesn't depend on l .

By the above theorem, we have when $0 < t \leq t_2$,

$$\sum_{n=0}^{\infty} \frac{\bar{q}_n(t, x, z)}{n!} \leq \sum_{n=0}^{\infty} \tilde{C}_1 K^n t^{-\frac{d}{\alpha}} \left(1 \wedge \frac{t^{\frac{1}{\alpha}}}{|x - z|} \right)^{d+\alpha} = \tilde{C}_1 \frac{1}{1 - K} t^{-\frac{d}{\alpha}} \left(1 \wedge \frac{t^{\frac{1}{\alpha}}}{|x - z|} \right)^{d+\alpha}.$$

Recall that $|q_n(t, x, z)| \leq \bar{q}_n(t, x, z)$, for any $n \geq 0$, i.e. $|q_n^{(l)}(t, x, z)| \leq \bar{q}_n(t, x, z)$, for any $n \geq 0$, because of the omission of the index l . It is clear that $\sum_{n=0}^{\infty} \frac{q_n^{(l)}(t, x, z)}{n!}$ is uniformly convergent on $[\epsilon, t_0] \times \mathbb{R}^d \times \mathbb{R}^d$, for any $\epsilon > 0$. Note that $\lim_{l \rightarrow \infty} q_n^{(l)}(t, x, z)$ exists, for any $n \geq 0$. We define

$q^{(l)}(t, x, z) = \sum_{n=0}^{\infty} \frac{q_n^{(l)}(t, x, z)}{n!}$. Then it is easy to see that $\lim_{l \rightarrow \infty} q^{(l)}(t, x, z)$ exists and its absolute value $\leq \bar{q}(t, x, z)$. We denote $\lim_{l \rightarrow \infty} q^{(l)}(t, x, z)$ by $q(t, x, z)$ and we have

$$|q(t, x, z)| \leq \tilde{C}_1 \frac{1}{1-K} t^{-\frac{d}{\alpha}} \left(1 \wedge \frac{t^{\frac{1}{\alpha}}}{|x-z|} \right)^{d+\alpha}. \quad (3.6)$$

Choosing l to be positive integers, we know that

$$A^{(l)}(t) = \sum_{s \leq t} F(X_{s-}, X_s) 1_{\{|X_{s-} - X_s| > 1/l\}} - \int_0^t \int_{\mathbb{R}^d} \frac{2C(X_s, y) F(X_s, y) 1_{\{|X_s - y| > 1/l\}}}{|X_s - y|^{d+\alpha}} m(dy) ds$$

is a local martingale. Define

$$L^{(l)}(t) = \exp\left(\sum_{s \leq t} (\ln(1 + F(X_{s-}, X_s) 1_{\{|X_{s-} - X_s| > 1/l\}})) - \int_0^t \int_{\mathbb{R}^d} \frac{2C(X_s, y) F(X_s, y) 1_{\{|X_s - y| > 1/l\}}}{|X_s - y|^{d+\alpha}} m(dy) ds\right),$$

and

$$L(t) = \exp\left(\lim_{l \rightarrow \infty} A^{(l)}(t) + \sum_{s \leq t} (\ln(1 + F) - F)(X_{s-}, X_s)\right).$$

By an argument similar to that used in [9], we have the following: the limit $\lim_{l \rightarrow \infty} A^{(l)}(t)$ exists in the sense of convergence of the norm of the space of square-integrable martingale and convergence in probability under P_x and the limit is also a martingale additive functional of X ; $L^{(l)}(t)$ converges to $L(t)$ in probability which is a local martingale and thus a supermartingale multiplicative functional of X ; $L(t)$ defines a family of probability of measures $\{\mathbb{P}_x, x \in \mathbb{R}^d\}$ on $\mathcal{M}_\infty$ by $d\tilde{\mathbb{P}}_x = L(t) d\mathbb{P}_x$ on $\mathcal{M}_t$. Let $\tilde{X} = (\tilde{X}_t, \mathcal{M}, \mathcal{M}_t, \tilde{\mathbb{P}}_x, x \in \mathbb{R}^d)$ denote this new process. Although $\tilde{X}_t(\omega) = X_t(\omega)$, we use $\tilde{X}_t$ to emphasize it is under $\tilde{\mathbb{P}}_x$. This process is a purely discontinuous Girsanov transform of X .

Since $L^{(l)}(t)$ converges to $L(t)$ in probability, there exists a subsequence $L^{(l_i)}(t)$ of $L^{(l)}(t)$ such that

$$\lim_{l_i \rightarrow \infty} L^{(l_i)}(t) = L(t).$$

On the other hand, since $\lim_{l \rightarrow \infty} q^{(l_i)}(t, x, z) = q(t, x, z)$ and $q(t, x, z)$ is integrable under the measure $m(dy)$ by (3.6), we have

$$\int_{\mathbb{R}^d} q(t, x, z) g(z) m(dz) = \mathbb{E}_x[L(t) g(X_t)],$$

for any bounded measurable function g , i.e. $q(t, x, z)$ is the transition density function of the purely discontinuous Girsanov transform of X . Thus the upper bound of the transition density function is obtained.

Next we establish the estimate of the lower bound of $q(t, x, z)$.

Lemma 3.4 *For any positive integer l , there exist $t_2 > 0$ and a large enough positive integer k such that when $0 < t \leq t_2$,*

$$\frac{|q_1^{(l)}(t, x, z)|}{k} \leq \frac{1}{2}p(t, x, z), \quad \forall (x, z) \in \mathbb{R}^d \times \mathbb{R}^d.$$

Proof. By Theorem 3.3, for any positive integer l , there exists $t_2 > 0$ which doesn't depend on l such that when $0 < t \leq t_2$,

$$|q_1^{(l)}(t, x, z)| \leq \bar{q}_1(t, x, z) \leq \tilde{C}_1 K t^{-\frac{d}{\alpha}} \left(1 \wedge \frac{t^{\frac{1}{\alpha}}}{|x - z|}\right)^{d+\alpha}.$$

On the other hand, by (1.1) we have

$$M_1 t^{-\frac{d}{\alpha}} \left(1 \wedge \frac{t^{\frac{1}{\alpha}}}{|x - z|}\right)^{d+\alpha} \leq p(t, x, z).$$

Thus it is easy to see that the statement holds. $\square$.

The above lemma implies that for any positive integer l , when $0 < t \leq t_2$,

$$p(t, x, z) + \frac{q_1^{(l)}(t, x, z)}{k} \geq p(t, x, z) - \frac{|q_1^{(l)}(t, x, z)|}{k} \geq \frac{1}{2}p(t, x, z).$$

We know $\int_{\mathbb{R}^d} \frac{q_1^{(l)}(t, x, z)}{k} m(dz) = \mathbb{E}_x[\frac{A_t^{(l)}}{k} g(X_t)]$, for any g bounded measurable. Let B_r be the ball $\{x \in \mathbb{R}^d : |x| \leq r\}$. Since $1 + \frac{A_t^{(l)}}{k} \leq e^{\frac{A_t^{(l)}}{k}}$, we have

$$\frac{1}{|B_r|} \mathbb{E}_x[(1 + \frac{A_t^{(l)}}{k}) 1_{B_r}(X_t)] \leq \frac{1}{|B_r|} \mathbb{E}_x[e^{\frac{A_t^{(l)}}{k}} 1_{B_r}(X_t)]$$

Thus

$$\begin{aligned} \frac{1}{2} \frac{1}{|B_r|} \mathbb{E}_x[1_{B_r}(X_t)] &\leq \frac{1}{|B_r|} \mathbb{E}_x[e^{\frac{A_t^{(l)}}{k}} 1_{B_r}(X_t)] \\ &\leq \left(\frac{1}{|B_r|} \mathbb{E}_x[e^{A_t^{(l)}} 1_{B_r}(X_t)]\right)^{\frac{1}{k}} \left(\frac{1}{|B_r|} \mathbb{E}_x[1_{B_r}(X_t)]\right)^{1-\frac{1}{k}} \\ &\quad (\text{by Hölder inequality}). \end{aligned}$$

Therefore

$$\begin{aligned} \frac{\frac{1}{2} \frac{1}{|B_r|} \mathbb{E}_x[1_{B_r}(X_t)]}{\left(\frac{1}{|B_r|} \mathbb{E}_x[1_{B_r}(X_t)]\right)^{1-\frac{1}{k}}} &\leq \left(\frac{1}{|B_r|} \mathbb{E}_x[e^{A_t^{(l)}} 1_{B_r}(X_t)]\right)^{\frac{1}{k}} \\ &\quad \text{i.e.} \\ \frac{1}{2} \left(\frac{1}{|B_r|} \mathbb{E}_x[1_{B_r}(X_t)]\right)^{\frac{1}{k}} &\leq \left(\frac{1}{|B_r|} \mathbb{E}_x[e^{A_t^{(l)}} 1_{B_r}(X_t)]\right)^{\frac{1}{k}} \\ &\quad \text{i.e.} \\ \frac{1}{2^k} \frac{1}{|B_r|} \mathbb{E}_x[1_{B_r}(X_t)] &\leq \frac{1}{|B_r|} \mathbb{E}_x[e^{A_t^{(l)}} 1_{B_r}(X_t)]. \end{aligned}$$

Let $r \downarrow 0$, we have when $0 < t \leq t_2$

$$\frac{1}{2^k} p(t, x, z) \leq q^{(l)}(t, x, z), \quad \forall l > 0.$$

Since $q(t, x, z) = \lim_{l \rightarrow \infty} q^{(l)}(t, x, z)$, we have

$$\frac{1}{2^k} p(t, x, z) \leq q(t, x, z).$$

Thus we obtain the lower bound of $q(t, x, z)$.

It is easy to see that the semigroup property of $e^{A_t^{(l)}}$ implies that

$$\int_{\mathbb{R}^d} q^{(l)}(t, x, y) q^{(l)}(s, y, z) dy = q^{(l)}(t + s, x, z), \quad \forall t, s > 0.$$

Thus by taking limit $\lim_{l \rightarrow \infty}$, we have

$$\int_{\mathbb{R}^d} q(t, x, y) q(s, y, z) dy = q(t + s, x, z), \quad \forall t, s > 0.$$

By this property and the lower and upper bounds of $q(t, x, z)$ when $0 < t \leq t_2$, we have the following theorem

Theorem 3.5 *There exist positive constants C_3, C_4, C_5 and C_6 such that*

$$C_3 e^{-C_4 t} t^{-\frac{d}{\alpha}} \left(1 \wedge \frac{t^{\frac{1}{\alpha}}}{|x - z|} \right)^{d+\alpha} \leq q(t, x, z) \leq C_5 e^{C_6 t} t^{-\frac{d}{\alpha}} \left(1 \wedge \frac{t^{\frac{1}{\alpha}}}{|x - z|} \right)^{d+\alpha} \quad (3.7)$$

for all $(t, x, z) \in (0, \infty) \times \mathbb{R}^d \times \mathbb{R}^d$.

Acknowledgement: I am very grateful to my advisor Professor Renming Song for his encouragement and many suggestions.

References

- [1] Ph. Blanchard and Z. M. Ma, Semigroup of Schrödinger operators with potentials given by Radon measures, in *Stochastic processes, physics and geometry*, 160-195, Word Sci. Publishing, Teaneck, NJ, 1990.
- [2] Z.-Q. Chen and R. Song, Conditional gauge theorem for non-local Feynman-Kac transforms, *Probab. Theory Relat. Fields*, **125**(2003), 45-72.
- [3] Z.-Q. Chen and R. Song, Drift transforms and Green function estimates for discontinuous processes, *J. Funct. Anal.*, 201(2003), 262-281.

- [4] Z.-Q. Chen and T. Kumagai, Heat kernel estimates for stable-like processes for stable-like processes on d -sets, *Stochastic Process. Appl.*, **108**(2003), 27-62
- [5] S. W. He, J. G. Wang and J. A. Yan, *Semimartingale Theory and Stochastic Calculus*, Science Press, Beijing New York, 1992.
- [6] M. Sharpe, *General Theory of Markov Processes*, Academic Press, Boston, 1988.
- [7] K. Sato, *Lévy processes and infinitely divisible distributions*, Cambridge University Press, Cambridge, 1999.
- [8] R. Song, Two-sides estimates on the density of the Feynman-Kac semigroups of stable-like processes, in preprint.
- [9] R. Song, Estimates on the Transition Densities of Girsanov Transforms of Symmetric Stable Processes, (to appear in *J. Theoret. Prob.*).
- [10] C. Wang, On the Estimates of the Density of the Feynman-Kac Semigroups of α -Stable-like Processes, in preprint.
- [11] Z. Vondraček, Basic potential theory of certain nonsymmetric strictly α -stable processes, *Glasnik Matematički.*, **37**(2002), 211-233.